

From: [Bridger, Michael C FLNR:EX](#)
To: [Scheck, Joelle L FLNR:EX](#)
Subject: RE: Pred mgmt
Date: Wednesday, June 5, 2019 2:07:31 PM
Attachments: [image001.png](#)
[image004.png](#)

Gotcha. I'll get this draft done by the end of the week. It will be a bit raw, as it's the very first draft prior to any sort of review. So maybe just highlight to COLT that it is the first draft. The results speak for themselves though.

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Scheck, Joelle L FLNR:EX
Sent: Wednesday, June 5, 2019 2:05 PM
To: Bridger, Michael C FLNR:EX
Subject: FW: Pred mgmt
Importance: High

Hi Mike,

See the string below. The issue that Vivian is raising doesn't apply to the Graham but she is looking for a 5 year review for Central Group to inform proposed predator management in SMC elsewhere. I have told Darcy and the rest of COLT that you are nearly done a first draft. If you have this by the end of this week (sorry, no pressure), then I will share it with COLT. I conveyed that we want to have some independent review of the review document before it is finalized.

Bevan will be assigned to undertake the issues note and summary note.

Joelle Scheck, RPBio, MScF
NE Caribou Team Lead
MFLNRORD - Regional Operations - Northeast
Suite 300, 10003 110th Avenue
Fort St. John, BC, V1J 5M9
Office: 250-787-3393
Cell: 250-261-1598



From: Peel, Darcy E FLNR:EX

Sent: June 5, 2019 12:07 PM

To: Wiebe, Heather I FLNR:EX <Heather.Wiebe@gov.bc.ca>; Scheck, Joelle L FLNR:EX <Joelle.Scheck@gov.bc.ca>

Subject: Fwd: Pred mgmt

Importance: High

FYI

Should have included you both in the distribution but hope the others can coordinate the development of the package

Darcy

Sent from my Samsung Galaxy smartphone.

----- Original message -----

From: "Peel, Darcy E FLNR:EX" <Darcy.Peel@gov.bc.ca>

Date: 2019-06-05 12:04 PM (GMT-08:00)

To: "Ernst, Bevan FLNR:EX" <Bevan.Ernst@gov.bc.ca>, "DeGroot, Leo FLNR:EX" <Leo.DeGroot@gov.bc.ca>, "Surgenor, John ENV:EX" <John.Surgenor@gov.bc.ca>

Subject: Pred mgmt

Hi guys, the comms folks are asking for some info re pred mgmt as well as directing us NOT start meetings until July.

Vivian wants a package with the following: 5 yr review and recommendations, issues note, summary note, and a Q&A

We have most of this either written already or in our collective heads. Can we get together asap to discuss how we address her concerns.

Thank you

Sent from my Samsung Galaxy smartphone.

Page 3 of 92
FNR 409-
96488

Page 4 of 92
FNR-2019-
96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Scheck, Joelle L FLNR:EX](#)
Subject: Draft South Peace Wolf Control Summary
Date: Friday, June 7, 2019 3:09:52 PM
Attachments: [image001.png](#)
[image003.png](#)

Here you go, Joelle. First draft attached. Let me know where you'd like to go from here with it. I'll send a copy to Bevan in the meantime as well.

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 6 of 92
FNR 409-
96488

Page 7 of 92
FNR-2019-
96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Ernst, Bevan FLNR:EX](#)
Subject: Draft South Peace Wolf Control Summary
Date: Friday, June 7, 2019 3:11:49 PM
Attachments: [image001.png](#)
[image003.png](#)

Hey Bevan,

I've attached the first draft of the 5-year summary report. Joelle also has a copy and will probably help decide a plan for getting this document reviewed. I'm guessing you'll be one of the people asked to review this eventually anyway.

Cheers,

Mike

Mike Bridger, M.Sc., R.P.Bio. 
Caribou Recovery Biologist | Northeast Region
Ministry of Forests, Lands, Natural Resource Operations and Rural Development
Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7
Ph: 250-787-3294



Page 9 of 92
FNR 409-
96488

Page 10 of
92 FNR-
2019-96488

From: [Grant, Laura FLNR:EX](#)
To: [Bridger, Michael C FLNR:EX](#)
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report
Date: Tuesday, July 2, 2019 10:32:56 AM
Attachments: [image001.png](#)
[image004.png](#)

No probs Bridge. It was a great (and very interesting) document, you just mentioned that no one else had looked at it. I know you don't have many folks up there to throw down solid edits, so I wanted to give you something to work with.

L

From: Bridger, Michael C FLNR:EX
Sent: July 2, 2019 10:29 AM
To: Grant, Laura FLNR:EX <Laura.Grant@gov.bc.ca>
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Thanks a lot Laura, these edits look really good. Sorry I kinda had to rush the document, so it is far from a polished product... so lots of edits required.

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Grant, Laura FLNR:EX
Sent: Friday, June 28, 2019 5:13 PM
To: Bridger, Michael C FLNR:EX
Cc: Roberts, Anne-Marie FLNR:EX; Surgenor, John ENV:EX; Ernst, Bevan FLNR:EX; Klaczek, Michael FLNR:EX; Roberts, Anne-Marie FLNR:EX; Scheck, Joelle L FLNR:EX
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hi Bridger,

Thank you for the opportunity to review this report. Very interesting results and you've done a great job of communicating the intricacies of a complex program! I have attached edits and comments. Please feel free to ignore any of them and get in touch if you have any questions or concerns.

Have a great weekend,

Laura

From: Roberts, Anne-Marie FLNR:EX
Sent: June 27, 2019 11:35 PM
To: Grant, Laura FLNR:EX <Laura.Grant@gov.bc.ca>
Subject: FW: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

FYI – perhaps you have seen this already but if not this is the report I mentioned today.

From: Bridger, Michael C FLNR:EX
Sent: June 26, 2019 11:53 AM
To: Surgenor, John ENV:EX <John.Surgenor@gov.bc.ca>; Ernst, Bevan FLNR:EX <Bevan.Ernst@gov.bc.ca>; Klaczek, Michael FLNR:EX <Michael.Klaczek@gov.bc.ca>; Roberts, Anne-Marie FLNR:EX <AnneMarie.Roberts@gov.bc.ca>
Cc: Scheck, Joelle L FLNR:EX <Joelle.Scheck@gov.bc.ca>
Subject: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hey all,

Some of you may have seen this document already, but I was hoping to get a proper review from a few folks from the Science Team and from each of the North Area regions. Would you be willing and able to give this document a read thru and provide edits and comments? It's very much a first draft at this point; no one other than myself has gone through it, so there is likely lots of rooms for edits and revisions. As it stands now, it is somewhat of an inward-facing document, as it discusses budgeting, contractor proficiency, etc. It would be good to take some of the key messages and create an outreach document as well.

After the internal review, our plan is to then send it out for an external review as well. We'll be asking the external reviewers to look at it from the lens of a potential scientific publication. We'll get one more year of caribou response data this coming winter to help strengthen the results.

Let me know if you think you'll have a bit of time to read through it. If not, no worries, just let me know and I'll check with some other folks. One part of the report I find a bit clunky is the year-by-year Results section. I think I could probably display that information in a table for each year and cut back on the text and the repetitiveness. Maybe let me know if you think that is a better way.

Thanks!

Mike

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 13 of
92
2019-96488

Page 14 of
92 FNR-
2019-96488

From: Klaczek, Michael FLNR:EX
To: Bridger, Michael C FLNR:EX
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report
Date: Friday, July 5, 2019 2:31:58 PM
Attachments: image001.png
image004.png

Awesome. Thanks. I've attached moose survey for Kennedy. We counted the Parsnip in 2017/18 – just south of your study area but similar density to pine/Mackenzie.

We did a comp in this pine/Mackenzie area in 2017/18 summarized in the Parsnip SRB report. Sex ratio was way up of calf:cow was similar.

s.22 . Shelley is around – Morgan/Matt trapping wolves in JPRF – I think still. They're there now anyway for a while.

From: Bridger, Michael C FLNR:EX
Sent: Friday, July 5, 2019 2:15 PM
To: Klaczek, Michael FLNR:EX
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Thanks man! Really good comments, I totally agree. I'll have a look through the primary prev discussion and see if I can emphasize your points. s.13

s.13

Are you in the office next week? I'm going to be in PG on Wednesdays. s.22

s.22 . Would be good to catch up with you and a few others at the office if you're around.

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Klaczek, Michael FLNR:EX
Sent: Thursday, July 4, 2019 6:17 PM
To: Bridger, Michael C FLNR:EX
Subject: Re: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hey Mike - I know the timelines are tight on this. I had a quick scan - looks awesome. s.13

s.13

s.13

s.13

I'll forward tomorrow.

Mike

From: Bridger, Michael C FLNR:EX
Sent: July-02-19 10:29 AM
To: Klaczek, Michael FLNR:EX
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Thanks man!

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Klaczek, Michael FLNR:EX
Sent: Friday, June 28, 2019 5:43 PM
To: Bridger, Michael C FLNR:EX
Subject: Re: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

I can review next week.

From: Bridger, Michael C FLNR:EX
Sent: June-26-19 11:52:56 AM
To: Surgenor, John ENV:EX; Ernst, Bevan FLNR:EX; Klaczek, Michael FLNR:EX; Roberts, Anne-Marie FLNR:EX
Cc: Scheck, Joelle L FLNR:EX
Subject: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hey all,

Some of you may have seen this document already, but I was hoping to get a proper review from a few folks from the Science Team and from each of the North Area regions. Would you be willing and able to give this document a read thru and provide edits and comments? It's very much a first draft at this point; no one other than myself has gone through it, so there is likely lots of room for edits and revisions. As it stands now, it is somewhat of an inward-facing document, as it discusses budgeting, contractor proficiency, etc. It would be good to take some of the key messages and create an outreach document as well.

After the internal review, our plan is to then send it out for an external review as well. We'll be asking the external reviewers to look at it from the lens of a potential scientific publication. We'll get one more year of caribou response data this coming winter to help strengthen the results.

Let me know if you think you'll have a bit of time to read through it. If not, no worries, just let me know and I'll check with some other folks. One part of the report I find a bit clunky is the year-by-year Results section. I think I could probably display that information in a table for each year and cut back on the text and the repetitiveness. Maybe let me know if you think that is a better way.

Thanks!

Mike

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 18 of
92
92-AR-
2019-96488

Page 19 of
92 FNR-
2019-96488

From: [Surgenor, John ENV:EX](#)
To: [Bridger, Michael C FLNR:EX](#)
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report
Date: Monday, July 8, 2019 1:48:20 PM
Attachments: [image001.png](#)
[image003.png](#)

Hi Mike, s.22 . Your report reads well. I don't really have any comments – that Laura hasn't touched on. Well done.

From: Bridger, Michael C FLNR:EX
Sent: July 5, 2019 4:43 PM
To: Surgenor, John ENV:EX <John.Surgenor@gov.bc.ca>; Ernst, Bevan FLNR:EX <Bevan.Ernst@gov.bc.ca>; Klaczek, Michael FLNR:EX <Michael.Klaczek@gov.bc.ca>; Roberts, Anne-Marie FLNR:EX <AnneMarie.Roberts@gov.bc.ca>; Grant, Laura FLNR:EX <Laura.Grant@gov.bc.ca>
Cc: Scheck, Joelle L FLNR:EX <Joelle.Scheck@gov.bc.ca>
Subject: RE: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hey there,

Laura put some serious effort into providing amazing edits and comments (got some good comments from Klaczek too), and I believe once I've finished incorporating those, the document will be much tighter and easier to read through. So for you folks, if you haven't dived deep into the edits already, I can re-send the document once I've finished working through Laura's work, and that will likely make things easier for you. I'm aiming to have those edits finished early next week, then I will re-send it to the group.

If you've already been working on it, all good, send it on over when you're done and I'll go through it.

Cheers!

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Bridger, Michael C FLNR:EX
Sent: Wednesday, June 26, 2019 11:53 AM
To: Surgenor, John ENV:EX; Ernst, Bevan FLNR:EX; Klaczek, Michael FLNR:EX; Roberts, Anne-Marie FLNR:EX
Cc: Scheck, Joelle L FLNR:EX
Subject: Internal Review of the 5-Year South Peace Pred. Mgmt. Summary report

Hey all,

Some of you may have seen this document already, but I was hoping to get a proper review from a few folks from the Science Team and from each of the North Area regions. Would you be willing

and able to give this document a read thru and provide edits and comments? It's very much a first draft at this point; no one other than myself has gone through it, so there is likely lots of rooms for edits and revisions. As it stands now, it is somewhat of an inward-facing document, as it discusses budgeting, contractor proficiency, etc. It would be good to take some of the key messages and create an outreach document as well.

After the internal review, our plan is to then send it out for an external review as well. We'll be asking the external reviewers to look at it from the lens of a potential scientific publication. We'll get one more year of caribou response data this coming winter to help strengthen the results.

Let me know if you think you'll have a bit of time to read through it. If not, no worries, just let me know and I'll check with some other folks. One part of the report I find a bit clunky is the year-by-year Results section. I think I could probably display that information in a table for each year and cut back on the text and the repetitiveness. Maybe let me know if you think that is a better way.

Thanks!

Mike

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 22 of
92
92-AR-
2019-96488

Page 23 of
92 FNR-
2019-96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Muter, David FLNR:EX](#)
Subject: RE: predator management plan
Date: Friday, July 12, 2019 1:34:27 PM

Thanks David.

Ya, fair point.^{s.13}
s.13
s.13 . That would be a good addition to the Introduction section at least.

Mike Bridger

Caribou Recovery Biologist

FLNRORD Northeast Region

Phone: (250)787-3294

From: Muter, David FLNR:EX
Sent: Friday, July 12, 2019 1:20 PM
To: Bridger, Michael C FLNR:EX
Subject: RE: predator management plan

Hi Mike,

This is very well done. Thanks for this.

I have one question - ^{s.13}
s.13

Thanks

D

<< OLE Object: Picture (Device Independent Bitmap) >> David Muter

Executive Director, Species at Risk Recovery

Resource Stewardship Division

(250) 217-5385

From: Bridger, Michael C FLNR:EX
Sent: July 8, 2019 4:59 PM
To: Muter, David FLNR:EX <David.Muter@gov.bc.ca>
Cc: Mitchell, Sean FLNR:EX <Sean.Mitchell@gov.bc.ca>; Ernst, Bevan FLNR:EX <Bevan.Ernst@gov.bc.ca>; Wiebe, Heather I FLNR:EX <Heather.Wiebe@gov.bc.ca>
Subject: RE: predator management plan

Hey David,

Got the latest edits finished. Here is the current draft version of the 5-Year report. Still not ready for distribution, but the content should be good to go.

Cheers,

Mike

<< File: Draft_South Peace Caribou Recovery - Wolf Control.docx >>

Mike Bridger

Caribou Recovery Biologist

FLNRORD Northeast Region

Phone: (250)787-3294

<< OLE Object: Picture (Device Independent Bitmap) >> << OLE Object: Picture (Device Independent Bitmap) >>

-----Original Appointment-----

From: Muter, David FLNR:EX
Sent: Friday, July 5, 2019 3:51 PM
To: Muter, David FLNR:EX; Mitchell, Sean FLNR:EX; Bridger, Michael C FLNR:EX; Ernst, Bevan FLNR:EX; Wiebe, Heather I FLNR:EX
Subject: predator management plan
When: Monday, July 8, 2019 11:30 AM-12:00 PM (UTC-08:00) Pacific Time (US & Canada).
Where: Skype Meeting

Hi all,

Wonder if we could connect to discuss next steps on predator management plan for this

year.

Thanks

D

Join Skype Meeting

Trouble Joining? [Try Skype Web App](#)

Join by phone

Local - Victoria: +1 (250) 952-9304, s.15;	(BC, Canada)	English (United States)
Local - Vancouver: +1 (604) 398-9304, s.15;	(BC, Canada)	English (United States)
Toll-Free: +1 (888) 952-9304, s.15;	(BC, Canada)	English (United States)

[Find a local number](#)

Conference ID: s.15;

[Forgot your dial-in PIN?](#) | [Help](#)

From: [Bridger, Michael C FLNR:EX](#)
To: [Marshall, Shelley FLNR:EX](#); [Klaczek, Michael FLNR:EX](#); [Jacklin, James FLNR:EX](#)
Subject: RE: bridger report
Date: Tuesday, July 16, 2019 1:42:14 PM
Attachments: [image001.png](#)
[image007.png](#)

Thanks Shelley, that's great!

I've incorporated or addressed most of those edits or comments.

A couple things... s.13
s.13

Good comments around the primary prey aspects. I've tightened up some of that wording and messaging.

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Marshall, Shelley FLNR:EX
Sent: Monday, July 15, 2019 8:23 PM
To: Bridger, Michael C FLNR:EX; Klaczek, Michael FLNR:EX; Jacklin, James FLNR:EX
Subject: Re: bridger report

This is super interesting and informative Mike. Because I just can't refrain, I've added some

questions and minor comments into the document despite not being asked to:-)

From: Bridger, Michael C FLNR:EX
Sent: July-15-19 10:16 AM
To: Klaczek, Michael FLNR:EX; Marshall, Shelley FLNR:EX; Jacklin, James FLNR:EX
Subject: RE: bridger report

Here you go. It's not quite ready for broad distribution just yet.

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Klaczek, Michael FLNR:EX
Sent: Monday, July 15, 2019 9:33 AM
To: Marshall, Shelley FLNR:EX; Jacklin, James FLNR:EX
Cc: Bridger, Michael C FLNR:EX
Subject: Re: bridger report

Hi Mike- do you have an updated draft you can forward to James? I only have the initial draft sent but I know there were comments/edits since.

Mike

From: Marshall, Shelley FLNR:EX
Sent: July-12-19 2:23 PM
To: Jacklin, James FLNR:EX; Klaczek, Michael FLNR:EX
Subject: RE: bridger report

It's not finalized yet I don't think. I believe Mike has reviewed a draft. I don't have a draft or final version.

From: Jacklin, James FLNR:EX
Sent: Friday, July 12, 2019 2:22 PM
To: Klaczek, Michael FLNR:EX
Cc: Marshall, Shelley FLNR:EX
Subject: bridger report

Can you please send me a copy of the 5 year pred mgt implementation report?

James Jacklin, R.P.Bio.
Director, Resource Management



Resource Management | Omineca Natural Resource Region | North Area
Phone (250) 561-3403
Forests, Lands, Natural Resource Operations and Rural Development

Page 30 of
92
2019-96488



Page 31 of 92

FNR-2019-06488

From: [Dale Seip](#)
To: [Bridger, Michael C FLNR:EX](#)
Subject: Re: 5-Year Wolf Reduction Report
Date: Wednesday, July 24, 2019 11:38:06 AM
Attachments: [image001.png](#)
[image003.png](#)

Hi Mike;

Report looks good. Just a few comments.

s.13;s.22

On Tue, Jul 16, 2019 at 2:22 PM Bridger, Michael C FLNR:EX
<Michael.Bridger@gov.bc.ca> wrote:

Hey^{s.22}

quick bit of work?

I've written a draft of the 5-Year Experimental Wolf Reduction for Central Mountain Caribou. It has been reviewed internally by a few folks, including some Caribou Science Team members. We're looking to have one or two folks provide an external review. I'm thinking this report would be of interest to you.

As it stands now, it's somewhat written as internal-facing, just because it does include some details around contractor proficiency, budgets, etc. After all of the final reviews, I'll probably create a version that is outward facing. The caribou program would like to be able to share the document broadly by late-August I believe.

So, would you be interested in reviewing it? It's about 30 pages, and I would send you the data as well. My timeline for receiving edits and comments would be around the 1st or 2nd week of August, because I'm taking off for a while starting the second half of August. We would just do a direct invoice with you, if you're able to provide an hourly or daily rate for your time.

One thing to keep in mind if you review it is the potential to turn this work into a scientific publication. I would like to get the 5th year of caribou response data first, just to increase the sample size. But would be curious to get your opinion if you think there would be enough good content for a publication. You come to mind as a potential co-author if we do decide to go down that road.

Let me know if you're interested and we can go from there.

Cheers!

MB

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 35 of
92
92-AR-
2019-96488

Page 36 of
92 FNR-
2019-96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Ernst, Bevan FLNR:EX](#)
Subject: South Peace Wolf Reduction Report
Date: Wednesday, August 7, 2019 11:25:02 AM
Attachments: [image001.png](#)
[image003.png](#)

Hey Bevan,

I think this document may be as good as it's going to get for now. It's been reviewed by Laura Grant, John Surgenor, yourself, Mike Klaczek, Shelley Marshall, Joelle Scheck, Dale Seip, and Rob Serrouya.

Do you want to start running it up the chain for the broader release? I've attached the Word and PDF versions.

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



Page 38 of
92
92-AR-
2019-96488

Page 39 of
92 FNR-
2019-96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Peel, Darcy E FLNR:EX](#)
Subject: RE: Predator Reduction for Caribou Recovery Consultation
Date: Monday, September 9, 2019 4:07:26 PM
Attachments: [image002.png](#)
[image005.png](#)

The cost per wolf is within that 5-year report for the South Peace. It's about \$5,100.

It was closer to \$2,500 in the Pink Mountain area. Really depends on several factors.

Mike Bridger
Wildlife Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Peel, Darcy E FLNR:EX
Sent: Friday, August 30, 2019 11:33 AM
To: Personal
Subject: RE: Predator Reduction for Caribou Recovery Consultation

Thanks ~~Pers~~ I will get you the information but it will likely be next week before I can pull it all together. I will also send you a copy of the 5 year review of the predator management program

Thank you for the phone call

Darcy

Darcy Peel – Director – Caribou Recovery Program
Cell: 778-220-3449



From: Personal Information
Sent: August 30, 2019 11:30 AM
To: Peel, Darcy E FLNR:EX <Darcy.Peel@gov.bc.ca>
Subject: Re: Predator Reduction for Caribou Recovery Consultation

As per our conversation regarding the cost of the wolf kills, could you please send me information as per cost per each dead wolf per hunt in the last two years using aircraft.

Thank you
Personal
Information

Sent from my iPad

On Aug 26, 2019, at 10:23 AM, "Peel, Darcy E FLNR:EX" <Darcy.Peel@gov.bc.ca> wrote:

Good morning,

On behalf of the Province of British Columbia, the Caribou Recovery Program is proposing an emergency predator reduction in the Tweedsmuir-Entiako, Hart Ranges, and Itcha-Ilgachuz herd areas.

This is a 30 day engagement process due to the urgency of the required action. See attached letter regarding engagement on this emergency management action. An information bulletin is also attached on the importance of wolf collars to caribou recovery.

For more information visit www.gov.bc.ca/caribou or to provide feedback on this consultation package, email – caribou.recovery@gov.bc.ca.

Thank you for your support and interest in Caribou

Darcy

Darcy Peel – Director – Caribou Recovery Program

Cell: 778-220-3449

<image001.png>

<predator reduction letter-all-final.pdf>

<Wolf Information Note-final.pdf>

Page 42 of
92
92-AR-
2019-96488



Page 43 of 92 FNR
-2019-96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Peel, Darcy E FLNR:EX](#); [Ernst, Bevan FLNR:EX](#)
Subject: RE: 5 year summary and FAQ
Date: Monday, September 23, 2019 8:32:41 AM
Attachments: [image001.png](#)
[image003.png](#)

Hey Darcy,

Here's the latest version of Bevan's annual summary report. It still says draft, but I think it's good to go... but may want to check with Bevan on that, I think he's back today. I know there is a good chance Sadie will post it publicly.

Mike Bridger
Wildlife Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Peel, Darcy E FLNR:EX
Sent: Saturday, September 21, 2019 1:41 PM
To: Ernst, Bevan FLNR:EX; Bridger, Michael C FLNR:EX
Subject: Fwd: 5 year summary and FAQ

Hi guys, I think you have already prepared this info but I don't have it handy. Could you forward info to me?

Thanks

Sent from my Samsung Galaxy smartphone.

----- Original message -----

From: Sadie Parr <sadie@wolfawareness.org>
Date: 2019-09-21 12:50 p.m. (GMT-08:00)
To: "Peel, Darcy E FLNR:EX" <Darcy.Peel@gov.bc.ca>
Subject: Re: 5 year summary and FAQ

Thank you for sending this report Darcy. How many wolves and other carnivores were killed in caribou ranges outside of the South Peace zone the last fiscal year? Is there an annual report from the past year that you can send that covers all areas where wolf killing is underway for caribou recovery?

Sincerely, Sadie Parr

On Thu, Sep 19, 2019 at 12:32 PM Peel, Darcy E FLNR:EX <Darcy.Peel@gov.bc.ca> wrote:

Hi Sadie, thank you for the conversation and I appreciate the level of concern you have. Attached you will find the 5 year summary as discussed as well as a FAQ document which addresses some of the key questions.

Darcy

Darcy Peel – Director – Caribou Recovery Program
Cell: 778-220-3449



Sadie Parr, Executive Director
Wolf Awareness Inc.

Research/Education/Conservation

T 250.272.HOWL (4695) E sadie@wolfawareness.org W wolfawareness.org + wehowl.ca

Page 46 of
92
92-AR-
2019-96488



Page 47 of 92 FNR
-2019-96488

From: [Bridger, Michael C FLNR:EX](#)
To: [Zimmerman, Kathi ENV:EX](#); [Dale.Seip@gmail.com](#); [Davis, Celine ENV:EX](#); [DeGroot, Leo FLNR:EX](#); [Dodd, Nicola L ENV:EX](#); [Doug Heard](#); [Ernst, Bevan FLNR:EX](#); [Friedrichsmeier, Christine FLNR:EX](#); [Grant, Laura FLNR:EX](#); [Klaczek, Michael FLNR:EX](#); [Pelletier, Agnes FLNR:EX](#); [Reid, Aaron FLNR:EX](#); [Roberts, Anne-Marie FLNR:EX](#); [Schildt, Scott FLNR:EX](#); [Schwantje, Helen FLNR:EX](#); [Shores, Carolyn FLNR:EX](#); [Thiessen, Conrad D FLNR:EX](#); [White, Shane FLNR:EX](#); [Williams, Bryan FLNR:EX](#); [Peel, Darcy E FLNR:EX](#); [Wiebe, Heather I FLNR:EX](#); [Burwash, Michael FLNR:EX](#); [Muhly, Tyler FLNR:EX](#); [Scheck, Joelle L FLNR:EX](#)
Subject: RE: F2F Science Team Meeting
Date: Wednesday, September 25, 2019 11:25:40 AM
Attachments: 2019 South Peace Caribou Recovery - Wolf Reduction.pdf

Apologies if this is a duplication for some of you, but I've attached the 5-Year wolf reduction summary report for the South Peace caribou recovery. Might be of value if you have a chance to flip through it prior to the meeting next week, as Bevan and I will be leading a bit of conversation around monitoring responses to wolf reduction, etc. I believe the report has been approved to be distributed externally, so feel free to pass it along if you like.

Mike Bridger

Wildlife Biologist

FLNRORD Northeast Region

Phone: (250)787-3294

-----Original Appointment-----

From: Zimmerman, Kathi ENV:EX

Sent: Tuesday, September 24, 2019 10:19 AM

To: Zimmerman, Kathi ENV:EX; Bridger, Michael C FLNR:EX; Dale.Seip@gmail.com; Davis, Celine ENV:EX; DeGroot, Leo FLNR:EX; Dodd, Nicola L ENV:EX; Doug Heard; Ernst, Bevan FLNR:EX; Friedrichsmeier, Christine FLNR:EX; Grant, Laura FLNR:EX; Klaczek, Michael FLNR:EX; Pelletier, Agnes FLNR:EX; Reid, Aaron FLNR:EX; Roberts, Anne-Marie FLNR:EX; Schildt, Scott FLNR:EX; Schwantje, Helen FLNR:EX; Shores, Carolyn FLNR:EX; Thiessen, Conrad D FLNR:EX; White, Shane FLNR:EX; Williams, Bryan FLNR:EX; Peel, Darcy E FLNR:EX; Wiebe, Heather I FLNR:EX; Burwash, Michael FLNR:EX; Muhly, Tyler FLNR:EX; Scheck, Joelle L FLNR:EX

Subject: F2F Science Team Meeting

When: Thursday, October 3, 2019 8:30 AM-4:00 PM (UTC-08:00) Pacific Time (US & Canada).

Where: 2975 Jutland Rd., 3rd Floor, Room W304

Good morning. Further to my placeholder email, this is the official invitation for the **Caribou Recovery SCIENCE TEAM Face to Face Meeting** in Victoria next Thursday, following the full program F2F meeting.

Attached please find the meeting agenda and objectives (I'll bring printed copies). This is an ambitious program to get through in one day, but I believe our previous 2-days of discussions will help set the stage for more detailed conversations about the role of the Science Team and the next steps we need to take.

Many thanks to those of you that have taken my calls and provided background and details on the topics you're working on, which has helped inform the agenda. Any additional agenda items you would like to propose on-site could be included in the final hour for those that do not need to leave early for travel.

Please take note that our meeting will kick-off with a round of introductions and I'd appreciate it if everyone could take 1-2 minutes to introduce yourself and present the top priority you are addressing and how the Science Team can support you.

Looking forward to seeing you next week!

Kathi

Kathi Zimmerman

Caribou Recovery Science Lead

Ministry of Environment

Kathi.Zimmerman@gov.bc.ca

M: 250-571-4751

P: 250-371-6282

<< File: Caribou Science Team Mtg Agenda - 2019-Oct-03.pdf >>



South Peace Caribou Recovery following Five Years of Experimental Wolf Reduction



Mike Bridger, R.P.Bio.

Wildlife Biologist | Northeast Region

BC Ministry of Forests, Lands, Natural Resource Operations and Rural Development

August 2019

Executive Summary

Experimental wolf reduction has occurred over a five-year period (2015-2019) within the South Peace region of northeastern British Columbia, Canada, in an attempt to address the rapid decline of Central Mountain caribou populations. These caribou herds have declined drastically in response to landscape changes that altered predator-prey dynamics and led to high rates of predation by wolves. The decrease in wolf abundance across the South Peace treatment area has shown conclusive evidence that intensive wolf reduction has halted and reversed the declining trends of the Klinse-Za, Kennedy Siding, and Quintette caribou populations. Although the first year of wolf reduction did not occur at a high enough intensity to elicit a caribou population effect, the following three years were sufficiently intensive (i.e. wolf densities were reduced to below 2 wolves/1000 km²) to elicit a strong, positive population response in all three treatment herds. The reduction of wolves during the fifth year is also expected to elicit a positive population response, but will not be measured until March 2020. As a result of wolf reduction, the South Peace caribou populations have increased by 49% from 166 individuals in 2016 to 247 individuals in 2019. The three herds combined had an average annual population growth rate of 15% following three effective years of wolf reductions and calf recruitment and adult female survival has increased in almost all cases in response to intensive wolf reductions. In contrast, prior to the implementation of wolf reduction, these three herds were declining at a rate of approximately 15% annually (625 to 166 individuals; 2002–2015). The adjacent, non-treatment caribou herds continued to show strong evidence of rapid declines over the same timeframe in the absence of wolf reduction.

Aerial wolf reduction has been shown to be the most targeted and effective method of intensively reducing wolf populations over large geographic areas to elicit strong population responses in caribou herds. Both the efficacy and efficiency of the South Peace wolf reduction program has increased over time. The success of the program is contingent on utilizing experienced and proficient removal crews, operating during optimal weather conditions, and maintaining a high level of operational oversight by provincial Ministry staff. Wolf reduction is a management tool that must be used responsibly and ethically, and implemented with the highest standards for humaneness and scientific rigour. Wolf reduction programs should be considered as an effective interim management tool for halting and reversing caribou declines, while the ultimate causes (i.e. habitat alteration) of such declines are addressed. Based on the findings of the five-year wolf reduction program in the South Peace, it is highly recommended that wolf reduction continue to be implemented to support these particular caribou herds towards meeting the ultimate management objective of self-sustaining populations.

Contents

Executive Summary.....	2
List of Tables.....	4
List of Figures.....	4
1. Introduction	5
2. Methods	7
3. Results	9
Wolf Reductions.....	9
Annual Caribou Population Results	10
Overall Caribou Population Results	11
Adjacent, Non-Treatment Herds.....	15
Wolf Recovery Rates.....	17
Primary Prey Response.....	19
4. Discussion.....	19
Caribou Population Results	19
Annual Wolf Recovery.....	20
Primary Prey	21
Links to Other Caribou Recovery Initiatives.....	22
Operational Considerations	22
5. Conclusion and Recommendations	24
6. Literature Cited.....	25

List of Tables

Table 1. Wolf reduction and caribou population results across all three South Peace caribou treatment herds in the Northeast Region of British Columbia, Canada.	12
--	----

List of Figures

Figure 1. South Peace herd boundaries for Central Mountain caribou, including local population unit (LPU) boundaries in the Northeast Region of British Columbia, Canada.	6
Figure 2. Treatment herds and wolf reduction boundaries across the South Peace caribou range in the Northeast Region of British Columbia, Canada.	8
Figure 3. Population trends for three South Peace caribou herds (Klinse-Za, Kennedy Siding, and Quintette) prior to, and in response to, intensive wolf reductions.	12
Figure 4. Average and overall population results for each treatment herd (green) in response to three years (2016-2018) of effective wolf reductions, and average population parameters for non-treatment caribou herds (red) over the same timeframe in the Northeast Region of British Columbia, Canada.	13
Figure 5. Annual caribou population growth rates relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The linear trend line intersects with a stable population growth rate at a density of approximately 7 wolves/1000 km ²	14
Figure 6. Proportion of caribou calves in the population relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The proportion of calves in the population that represents population stability is approximately 15.5% (Bergerud 1992).	14
Figure 7. Adult female caribou survival rates relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The adult female survival rate threshold for population stability is approximately 88% (Bergerud and Elliot 1986).	15
Figure 8. Proportion of calves in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment populations (Graham herd and South Narraway herd).	16
Figure 9. Adult female survival rate observed in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment population (Graham herd). ...	16
Figure 10. Annual caribou population growth rates observed in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment population (Graham herd). Lambda was calculated in the Graham population using Hatter and Bergerud’s (1991) recruitment-mortality equation.	17
Figure 11. Wolf recovery rates relative to the initial population of wolves (n = 208) in the treatment area.	18
Figure 12. Annual wolf recovery rates following wolf reductions across the treatment area, with a threshold of greater than 85% indicating reduced annual recovery rates.	18

1. Introduction

Throughout the 1990s and early 2000s, woodland caribou (*Rangifer tarandus caribou*) herds in the South Peace region of northeast British Columbia (BC) were presumed to be declining (Seip and Jones 2014). Increased monitoring efforts through the 2000s confirmed that these caribou herds were in fact decreasing at a rapid rate (Seip and Jones 2016). The status of the Central Mountain Designatable Unit (DU8; COSEWIC 2011) of woodland caribou found in the South Peace (Figure 1) has recently been updated to “Endangered” by the Committee on the Status of Endangered Wildlife in Canada. The South Peace herds include the Scott East and Moberly (which were combined in 2015 and are hereafter referred to as the Klinse-Za herd), Burnt Pine, Kennedy Siding, Quintette, Bearhole-Redwillow, and South Narraway (transboundary with Alberta). Within the South Peace region, these herds have further been grouped by local population units (LPUs), which includes the Pine LPU (composed of the Klinse-Za, Kennedy Siding, and former Burnt Pine herds), the Quintette LPU (composed of the Quintette herd), and the Narraway LPU (composed of the Bearhole-Redwillow and South Narraway herds). Prior to wolf reduction, all of these herds had been declining drastically, and the Burnt Pine herd was extirpated. These drastic declines followed extensive landscape change resulting from forest harvest, mining, oil and gas exploration, road construction and other industrial activities within or adjacent to caribou ranges. This has led to the direct loss of habitat and altered predator-prey dynamics. Many industrial activities promote early seral forests, which benefit species like moose (*Alces americanus*) and ultimately lead to increased wolf (*Canis lupus*) populations (moose are a primary prey species for wolves; Fuller et al. 2003). Such increases in wolf populations result in higher caribou mortality rates (Seip 1992), which can be further exacerbated by newly developed linear features that enhance wolf movement and provide access into caribou range. Wolf predation in the South Peace was occurring at rates that were unsustainable for caribou populations, leading to rapid population declines (Seip and Jones 2014).

Amongst the Central Mountain caribou herds found in the South Peace region, annual mortality rates of radio-collared adult females ranged from 12–24% (Seip and Jones 2014) prior to wolf reduction. Wolf predation accounted for 38% of all documented caribou mortalities, and 78% of all cases in which a conclusive cause of mortality was determined (Seip and Jones 2014). Calf recruitment ranged between 9–14% calves within the population (measured annually in late-March; Seip and Jones 2016), which was generally inadequate to compensate for adult mortality. The causes of calf mortality have not been investigated across the South Peace caribou herds; however, studies have shown wolves to be a significant predator of caribou calves in other jurisdictions (Gustine et al. 2006), and calf survival has been shown to increase in response to wolf reductions (Farnell and McDonald 1988, Seip 1992, Bergerud and Elliot 1998, Hayes et al. 2003). Previous research by Bergerud and Elliot (1986) concluded that wolf densities greater than 6.5 wolves/1000 km² resulted in caribou population declines. Furthermore, the Federal government’s recovery strategy for woodland caribou recommends a target wolf density of less than 3 wolves/1000 km² across caribou range (Environment Canada 2014). Prior to the implementation of wolf reduction, wolf densities across the South Peace caribou herds were estimated at approximately 10–14 wolves/1000 km² (Seip and Jones 2014); well above the density thresholds associated with the persistence of caribou populations. The pre-reduction wolf density estimate has since been refined to 12.6 wolves/1000 km².

In response to dramatic caribou population declines, provincial wildlife managers from the BC Ministry of Environment and Ministry of Forests, Lands, Natural Resource Operations and Rural Development (hereafter referred to as the Province or Ministry) approved the implementation of predator management in the form of aerial wolf reduction. The initial approval was for a five-year aerial wolf reduction program, which commenced during the winter of 2014-2015, and has since completed its fifth year of reductions following the winter of

2018-2019. The wolf reduction program has occurred in combination with other recovery efforts, including maternal penning, supplemental feeding, and habitat restoration. In 2017-2018, the program was expanded to include the South Narraway caribou herd range. Wolf reduction was implemented as an interim management measure to assist the South Peace caribou herds in reaching self-sustaining status, with a population target of approximately 1,000 individuals (800 combined in the Pine and Quintette LPUs and 200 in the Narraway LPU). Serrouya et al. (2019) reported similar conclusions regarding population growth rates of the South Peace treatment herds; however, the following report has been developed to further investigate the mechanisms and drivers of caribou population change, such as variation in calf recruitment and adult survival, relative to wolf reduction treatments.




Figure 1. South Peace herd boundaries for Central Mountain caribou, including local population unit (LPU) boundaries in the Northeast Region of British Columbia, Canada.

2. Methods

Prior to the implementation of wolf reductions, baseline population data for both caribou and wolves were collected. These data were collected continually over the five-year study period to measure caribou population responses. Caribou population data, including population estimates and calf recruitment rates, were measured annually through aerial surveys. Adult female survival rates were calculated by monitoring a subsample of radio-collared individuals within the population, and cause of mortality was determined through mortality investigations of deceased radio-collared individuals. Wolf pack locations, habitat use, and density estimates were derived through radio-collaring of wolves. Additionally, the response by primary prey species (i.e. moose) to wolf reductions was also measured within and adjacent to the treatment areas through aerial surveys and radio-collaring studies.

The initial treatment area boundary was designed to encompass the majority of the Klinse-Za, Kennedy Siding, and Quintette caribou herd ranges; a wolf reduction zone of approximately 16,500 km² (Figure 2). Specifically, the boundary included the core high elevation and low elevation caribou habitat, and adjacent matrix habitat within the caribou ranges. The South Narraway treatment area was included in 2017-2018, with an additional area of approximately 1,600 km². These treatment areas formed the boundaries for intensive reduction of wolves using aerial gunning from helicopters, where the objective was to remove the majority of wolves within or immediately adjacent to the reduction zone and in doing so, reduce wolf densities to below 3 wolves/1000 km². Aerial gunning of wolves was deemed the most effective and humane method of removal, as properly applied shooting techniques results in wolves being quickly dispatched while eliminating the risk of bycatch. The reduction of wolves occurred during the winter months when snow levels facilitated optimal tracking conditions, and concentrated wolves' distribution at lower elevations. Reduction efficiency was increased by deploying radio-collars on individual wolves in all known wolf packs within or immediately adjacent to the treatment area boundary. This facilitated the relocation of the wolf packs and increased the likelihood of removing all wolves from each pack. The individual wolves were captured via helicopter net-gunning, which enabled crews to restrain the individuals and deploy GPS-satellite radio-collars, allowing for remote tracking of movements and locations, and relocation through the use of radio telemetry. The radio-collared individuals were often left alive following the conclusion of the winter reduction efforts in order to facilitate the location of wolves the following winter. Wolves that were found immediately adjacent to the reduction zone, or were tracked from within the boundary to adjacent areas were also removed (assuming these wolves had at least partial overlap with the treatment area).

Aerial wolf reduction was delivered primarily by external contractors, with operational oversight from Ministry staff. Overtime, the operational oversight was increased in order to ensure the efficacy and humaneness of the program and internal Ministry staff assisted with the delivery of the field operations as well. Initially, wolf removal crews attempted to retrieve the carcasses of deceased wolves; however, it was quickly determined to be an inefficient use of time, effort, and funds. Subsequently, the locations of accessible wolf carcasses were provided to First Nations and they retrieved those carcasses from the ground. The Province also collaborated with local First Nations to support wolf reduction through ground trapping programs. Although deemed ineffective on its own (Webb et al. 2011), ground trapping was thought to offer an additional source of wolf removal, while providing opportunities to collaborate with local First Nations communities. The ground

trapping efforts were generally focused within the Klinse-Za caribou range, in close proximity to local First Nations communities and the caribou maternal penning site¹.




Figure 2. Treatment herds and wolf reduction boundaries across the South Peace caribou range in the Northeast Region of British Columbia, Canada.

¹ Maternal penning was an ongoing management initiative in the Klinse-Za caribou range that was occurring concurrently with wolf reduction (McNay et al. 2019)

The reduction of wolves occurred during five winters (2015-2019) and the response within caribou populations was measured during each of the following winters. Aerial surveys of most herds were conducted annually to estimate population size, calf survival/recruitment, and calculate population growth rates (also referred to as λ ; based on annual changes in population estimates). The Kennedy Siding herd was monitored via motion-sensitive cameras at supplemental feeding sites², allowing researchers to obtain total counts of the population, including population demographics (Heard and Zimmerman 2018). Additionally, a subsample of radio-collared female caribou was maintained in each herd during the five-year program to calculate adult female survival rates. Two adjacent, non-treatment herds (Graham and South Narraway herds) were identified as the experimental control populations for this program, allowing for population and demographic parameters to be compared between treatment and non-treatment herds over time. The South Narraway herd was subsequently removed as a non-treatment herd in 2017-2018 due to its continued rapid decline and urgent need for recovery measures (i.e. wolf reduction).

An initial wolf density estimate within the focal caribou herd ranges was derived through the radio-collaring of the majority of wolf packs in the area and through comparisons of previously recorded wolf densities observed in similar mountainous caribou ranges in neighbouring jurisdictions (Kuzyk 2002, Hayes et al. 2003). The estimates were further refined over the course of each winter's reduction efforts by documenting all wolves and wolf packs encountered during wolf removal flights. This allowed for relatively accurate documentation of the proportion of wolves removed each winter, and the subsequent density of wolves remaining following each winter's reduction efforts. The wolf density estimates were also calculated at the LPU level based on the number of wolves removed and number remaining following each winter season of wolf reduction. Similarly, wolf densities were also estimated across the non-treatment caribou herd ranges using a combination of radio collaring and extrapolation (i.e. Graham and South Narraway herds).

3. Results

Wolf Reductions

The wolf reduction program to support caribou recovery in the Klinse-Za, Kennedy Siding, and Quintette herds was initially approved during the winter of 2014-2015. The late approval for the program resulted in a delayed start to the field operations and an underspending of the budget (approximate cost of \$200,000). This late start, combined with poor winter conditions (i.e. lack of snow) and relatively new removal crews, led to an ineffective wolf reduction effort. The initial wolf population estimate within or immediately adjacent to the treatment area was 208 wolves and a density of 12.6 wolves/1000 km². Overall, 57 wolves were removed from the reduction zone (41 removed by aerial gunning, 16 removed by ground trapping), equating to a reduction of only 27% of the wolf population. There was an estimated 151 wolves remaining in the treatment area, and a density of 9.4 wolves/1000 km² following the wolf reduction efforts. The remaining density estimates at the LPU scale were 10.8 wolves/1000 km² in the Pine LPU, and 6.5 wolves/1000 km² in the Quintette LPU.

Three external contractors were hired to deliver the majority of the field operations for the second year (2015-2016) of the wolf reduction program, with minimal operational oversight or field involvement from Ministry staff. Overall, 201 wolves were removed across the treatment area (155 removed by aerial gunning, 46 removed by ground trapping). The reduction rate was estimated at 97%, with only seven wolves remaining in the treatment area and a density estimate of 0.4 wolves/1000 km² following reduction efforts. At the LPU level, there was a remaining density estimate of 0.5 wolves/1000 km² in the Pine LPU, and 0.3 wolves/1000 km² in the

² Supplemental feeding is an ongoing management initiative in the Kennedy Siding caribou range (Heard and Zimmermann 2018)

Quintette LPU. Although the level of wolf reduction during the second year was very high, a lack of Ministry oversight resulted in inflated program costs of approximately \$800,000.

The program was primarily delivered by two external contractors in the third year (2016-2017) of the program. There was a slight increase in operational oversight by Ministry staff, but minimal field involvement. Upon the conclusion of the third year of reduction efforts, an adequate number of wolves had been removed at a cost of approximately \$475,000. Overall, 103 wolves were removed (62 by aerial gunning, 31 by ground trapping, and an additional 10 by ground shooting at den sites in the spring). The wolf reduction rate was estimated at 79%, with 27 wolves remaining in or adjacent to the treatment area and a density of 1.7 wolves/1000 km² following the reduction efforts. Within the Pine LPU, there was an estimated density of 2.0 wolves/1000 km² remaining, and 1.1 wolves/1000 km² remaining in the Quintette LPU.

The fourth year (2017-2018) of wolf reductions was delivered by one primary contractor and a secondary crew led by Ministry staff. The level of operational oversight by Ministry staff was increased significantly. The wolf reduction efforts upon conclusion of the winter season were highly successful, both in terms of reduction efficacy and cost efficiency (total cost of \$376,000). Overall, 116 wolves were removed across the three treatment herds (all by aerial gunning; there were no conclusive reports of ground trapping removal). The wolf reduction rate was estimated at 92%, with only 10 wolves remaining within or immediately adjacent to the treatment area following the reduction efforts. This equated to a remaining density of 0.6 wolves/1000 km². At the LPU level, there was a density estimate of 0.3 wolves/1000 km² in the Pine LPU, and 1.1 wolves/1000 km² in the Quintette LPU. The winter of 2017-2018 also marked the first year of wolf reduction in the South Narraway caribou range. Fourteen wolves were removed on the BC side of the border (an additional 10 wolves were removed on the Alberta side). The reduction efforts in BC equated to an estimated reduction rate of 74%, with approximately five wolves remaining and a density of 3.1 wolves/1000 km². The total cost of conducting wolf reductions in the South Narraway was \$81,000.

During the fifth year (2018-2019) of wolf reductions, field operations were delivered primarily by one contractor, with a high level of operational oversight from Ministry staff. The wolf reduction efforts were once again effective and relatively efficient, with a total cost of approximately \$340,000. Overall, 61 wolves were removed (51 by aerial gunning, 10 by ground trapping). This equated to a reduction rate of 77%, with an estimated 18 wolves remaining within or immediately adjacent to the treatment area and a density of 1.1 wolves/1000 km². There was a remaining wolf density estimate of 1.0 wolves/1000 km² in the Pine LPU, and 1.3 wolves/1000 km² in the Quintette LPU. Within the South Narraway caribou range, it appeared that the wolf recovery rate was extremely low following the previous winter's reduction efforts. Only one wolf was removed from the BC side of the South Narraway range in 2018-2019, resulting in a remaining wolf density estimate of 1.9 wolves/1000 km².

Annual Caribou Population Results

The level of wolf reduction during the first year of wolf removals did not lead to a positive caribou population response, as evidenced by the population parameters reported the following year. Across the three caribou herds, the population had declined by an additional 13.5% ($\lambda = 0.865$) over the course of the year, the adult female survival rate was 78.7% ($n = 41$), and the proportion of calves in the population was 15.6% (measured late-March 2016). Within the Pine LPU, the population had increased by 14% ($\lambda = 1.14$), with an adult female survival rate of 85.7% ($n = 24$), but only 13.0% calves in the population. The Quintette LPU had declined by 38.0% ($\lambda = 0.62$), adult female survival was 64.7% ($n = 17$), however there were 20.0% calves in the population.

Following the second year of wolf reductions, the caribou population had increased across all three herds by 15.6% ($\lambda = 1.156$), the adult female survival was 92.7% ($n = 41$), and the proportion of calves in the population was 22.9%. Within the Pine LPU, the population had increased by 19.5% ($\lambda = 1.195$), with an adult female survival rate of 93.1% ($n = 29$) and 24.5% calves in the population. The Quintette LPU had increased by 10% ($\lambda = 1.10$), the adult female survival was 91.6% ($n = 12$) and 18.4% calves in the population.

Following the third year of wolf reduction, the caribou population increased by 6.7% ($\lambda = 1.067$), the adult female survival rate was 91.8% ($n = 46$), and the proportion of calves in the population was 18.6% (measured late-March 2018). Within the Pine LPU, the population had increased by 5.5% ($\lambda = 1.055$), with an adult female survival rate of 87.1% ($n = 31$) and 16.8% calves in the population. Within the Quintette LPU, the population increased by 9.0% ($\lambda = 1.09$), the adult female survival was 100% ($n = 15$), and 19.4% calves in the population.

The fourth year of wolf reductions led to a caribou population increase of 22.0% ($\lambda = 1.22$), the adult female survival rate was 89.2% ($n = 46$), and the proportion of calves in the population was 21.0% (measured late-March 2019). Within the Pine LPU, the population increased by 23.5% ($\lambda = 1.235$), with an adult female survival rate of 92.6% ($n = 27$), and 19.5% calves in the population. Within the Quintette LPU, the population increased by 18.9% ($\lambda = 1.189$), with an adult female survival rate of 77.8% ($n = 9$), and 25.0% calves in the population. Aerial survey efforts in the South Narraway documented a minimum observation of 38 caribou (up from 26 in March 2018), the adult female survival was 100% ($n = 12$), but only 13.2% calves in the population. The increase in caribou observations was likely explained by differences in survey efficacy between years.

Based on the results from previous years, the level of wolf reduction achieved during the fifth year of the program should be sufficient to elicit a strong, positive response in the caribou populations. The results of these reduction efforts, however, will not be measured until March 2020. According to the documented population trends, it is predicted that the fifth year of wolf reduction efforts will elicit approximately 15% caribou population growth ($\lambda = 1.15$), 90% adult female survival, and 21% calves in the population, across all three South Peace treatment herds. The effects of wolf reductions towards the South Narraway herd, and subsequent predictions, will require further investigation and monitoring.

Overall Caribou Population Results

The overall results measured during the five-year wolf reduction program suggest that the reduction of wolves to low densities can have significant, positive effects towards caribou populations. The level of wolf reduction during the first year did not lead to a caribou population response, however, the following three years of wolf reduction resulted in positive caribou responses in almost all population parameters measured (i.e. λ , adult female survival, and calf recruitment) across all treatment herds (Table 1). The level of wolf reduction achieved in Year 5 is expected to elicit similar responses, but will not be measured until March 2020.


The total population size across the three treatment herds had increased from 166 individuals in 2016 to 247 individuals in 2019 (a 49% population increase). The average annual population growth rate following the three years of effective wolf reduction was 15% ($\lambda = 1.15$). The Pine LPU increased from 104 individuals to 159, with a total population increase of 53%. The Quintette LPU increased from 62 individuals to 88, with a total population increase of 42%. When forecasting the future population trend, assuming an average annual growth rate of 15% and considering density-dependent growth, the caribou population across the three treatment herds could double in size by year 2027 and approach the population objective ($n = 800$) by 2037 (Figure 3).

Table 1. Wolf reduction and caribou population results across all three South Peace caribou treatment herds in the Northeast Region of British Columbia, Canada.

	2015	2016	2017	2018	2019	2020
Caribou population estimate	192*	166	192	205	247	275**
Calf recruitment	15.5%*	15.6%	22.9%	18.6%	21.0%	21.0%**
Adult female survival	82.5%*	78.7%	92.7%	91.8%	89.2%	90.0%**
Annual population growth rate	-10.0%*	-14.0%	15.6%	6.7%	22.0%	15.0%**
Proportion of wolves reduced during previous winter	0%*	27%	97%	79%	92%	77%
Individual wolves remaining after previous winter's reduction	208*	151	7	27	10	18
Wolf density remaining after previous winter's reduction (wolves/1000 km ²)	12.6*	9.4	0.4	1.7	0.6	1.1

*Parameters measured prior to the implementation of wolf reduction

**Predicted caribou population response based on 2018-2019 wolf reduction

**Figure 3.** Population trends for three South Peace caribou herds (Klinse-Za, Kennedy Siding, and Quintette) prior to, and in response to, intensive wolf reductions.




Figure 4. Average and overall population results for each treatment herd (green) in response to three years (2016-2018) of effective wolf reductions, and average population parameters for non-treatment caribou herds (red) over the same timeframe in the Northeast Region of British Columbia, Canada.

Prior to implementing wolf reductions (wolf density of 12.6 wolves/1000 km²) and following an ineffective wolf reduction effort (Year 1 – wolf density of 9.4 wolves/1000 km²), all measured caribou population parameters were indicative of declining populations. In each year following effective wolf reduction (i.e., reduction to two wolves/1000 km² or less), the population growth rate (Figure 5), calf recruitment (Figure 6), and adult female survival (Figure 7) were indicative of increasing caribou populations in all cases. The linear trend relating caribou population growth to wolf density appeared to suggest stable caribou populations could exist at a wolf density of approximately seven wolves/1000 km² (similar to the wolf density equilibrium reported by Bergerud and Elliot [1986]).




Figure 5. Annual caribou population growth rates relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The linear trend line intersects with a stable population growth rate at a density of approximately 7 wolves/1000 km².




Figure 6. Proportion of caribou calves in the population relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The proportion of calves in the population that represents population stability is approximately 15.5% (Bergerud 1992).




Figure 7. Adult female caribou survival rates relative to wolf density following reduction efforts across the three treatment herds. The “star” symbol represents the predicted response based on wolf reduction following the winter of 2018-2019. The adult female survival rate threshold for population stability is approximately 88% (Bergerud and Elliot 1986).

Adjacent, Non-Treatment Herds

During the years of wolf reduction across the South Peace caribou ranges, the adjacent non-treatment herds (i.e. Graham and South Narraway herds) showed demographic parameters suggestive of continued population declines. The wolf densities across the non-treatment herd ranges were assumed to have remained constant during that time, with an estimate of 12–14 wolves/1000 km² in the Graham caribou range and 11–12 wolves/1000 km² in the South Narraway core caribou range (FLNRORD *unpubl.*). During the years of effective wolf reduction in the treatment herds (2016-2019), the proportion of calves in the Graham herd ranged between 11.6–16.2% (compared to 16.8–25.0% in treatment herds; Figure 8). The adult female survival rate during that timeframe ranged between 60.0–80.8% (compared to 77.8–100% in treatment herds; Figure 9). The calf recruitment rate in the Graham herd was insufficient to compensate for the adult mortality rate, indicating a declining population trend. Calf recruitment in the South Narraway herd had ranged between 13.2–17.4%; however, there was an insufficient sample size of radio-collared caribou to monitor adult survival rates. The annual population growth rate for the Graham herd³ had continued to suggest negative growth, declining between 10.3–31.9% annually (Figure 10), while the treatment herds exhibited positive population growth in all years following effective wolf reductions (5.5–23.5% population growth).

³ Population estimates via aerial census had not been derived consistently for the Graham caribou herd. Lambda (λ) was calculated using Hatter and Bergeruds' (1991) recruitment-mortality equation, where: $\lambda = \text{adult female survival rate} / (1 - \text{recruitment})$




Figure 8. Proportion of calves in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment populations (Graham herd and South Narraway herd).




Figure 9. Adult female survival rate observed in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment population (Graham herd).




Figure 10. Annual caribou population growth rates observed in the treatment populations (Pine LPU and Quintette LPU) following three effective years of wolf reductions compared to the non-treatment population (Graham herd). Lambda was calculated in the Graham population using Hatter and Bergerud's (1991) recruitment-mortality equation.

Wolf Recovery Rates

The rate at which wolves recovered⁴ in the treatment area relative to the initial population varied depending on the proportion of wolves that were removed during previous winters (Figure 11). Following wolf reduction in Year 1, when only 27% of wolves were removed, there appeared to be 100% recovery of wolves relative to the initial population. After removing approximately 97% of the wolves in Year 2, wolves recovered at a rate of 63% of the initial population the following winter, and a similar rate was observed following Year 3 (61% recovery relative to the initial population). After achieving a 92% wolf reduction in Year 4, wolves appeared to have recovered to only 38% of the initial population the following winter. The recovery rate after Year 5 of wolf reductions will be measured during the winter of 2019-2020. Over time, the results suggest that wolf recovery rates can be reduced with successive years of intensive reduction efforts.

Furthermore, the level of wolf reduction necessary to reduce wolf recovery rates on an annual basis appeared quite high (Figure 12). For example, the removal of 79% of wolves in Year 3 still resulted in a high annual recovery rate of approximately 97%. Only following winters when wolves were reduced by greater than 90% (Years 2 and 4) were annual wolf recovery rates reduced to approximately 63%. The preliminary results suggest that a reduction of over 85% of wolves each winter is required to reduce annual recovery rates.

⁴ Either through population growth (i.e. producing litters) or recolonization (which occurs between the timeframe following the conclusion of wolf reduction at the end of the winter and commencement of wolf reduction at the beginning of the following winter)




Figure 11. Wolf recovery rates relative to the initial population of wolves (n = 208) in the treatment area.




Figure 12. Annual wolf recovery rates following wolf reductions across the treatment area, with a threshold of greater than 85% indicating reduced annual recovery rates.

Primary Prey Response

Primary prey species, particularly moose, were expected to respond to wolf reductions at a level equal to, or greater than the response observed in the caribou populations. However, baseline data for moose populations across the entire treatment area were incomplete, making for difficult interpretation of population trends. Moose populations have been monitored more frequently in recent years since the implementation of wolf reductions. During the winter of 2016-2017, an aerial population composition survey for moose in the Quintette caribou range reported a ratio of 38 calf moose per 100 cow moose (FLNRORD *unpubl.*). This result does not suggest a rapidly increasing moose population (more likely stable to increasing); however, this result occurred following only one year of effective wolf reductions, thus additional data were required to inform population response. During the winter of 2018-2019, a comprehensive moose survey occurred within the Quintette caribou range and yielded a moose density estimate of 0.19 moose/km² and a ratio of 33 calves per 100 cows (FLNRORD *unpubl.*). Despite three years of intensive wolf reduction prior to the survey, the moose population density was relatively low with a calf ratio suggesting a stable population. Within the Kennedy Siding herd range, a moose inventory was conducted in 2015, just prior to the implementation of wolf reduction, but has not been surveyed since. At the time of the survey, there was a density estimate of 0.49 moose/km² and 36 calves per 100 cows (Klaczek et al. 2016).

Within the Klinse-Za caribou range, a moose survey in 2017-2018 (following two years of effective wolf reduction) reported a moose density estimate of 0.44 moose/km² (no significant change from the previous estimate in 2012-2013) and a calf ratio of only 21 calves per 100 cows (Sittler and McNay 2018). Additionally, a comprehensive moose survival study in the Klinse-Za caribou range had occurred concurrently with wolf reduction efforts, and compared moose population parameters in response to wolf reductions with those of an adjacent, non-treatment population of moose (Sittler 2019). Overall, adult female moose had a 7% greater survival rate in the Klinse-Za range than in the non-treatment moose population, including one year of 100% survival (n = 40) following Year 4 of wolf reduction (when 96% of wolves in the Klinse-Za range were removed). Adult female survival ranged from 92–100% following three years of effective wolf reduction, whereas survival in the non-treatment population ranged from 83–93%. Calf recruitment was lower than expected in two of the years following effective wolf reductions, with a higher calf recruitment rate observed in the non-treatment population. Calf recruitment in the treatment population has since increased to 35 calves per 100 cows in March 2019. Lambda calculations suggested positive population growth rates in all years following wolf reduction (as was observed in the non-treatment population). Over the course of the study, wolf predation was the cause of one of five (20%) moose mortalities in the Klinse-Za range, while wolf predation was the cause of four of 11 mortalities (36%) in the non-treatment population.

4. Discussion

Caribou Population Results

The positive response measured across the South Peace caribou population following intensive wolf reduction provides strong support for the use of wolf reduction as a tool to increase caribou herds. The results show a conclusive, positive response by the treatment herds following three years of effective wolf reductions, whereas non-treatment caribou herds continued to show evidence of rapid declines. It was evident that a high level of wolf reduction was required to elicit the positive response in treated caribou populations; reduction of wolf densities to < 7 wolves/1000 km² may contribute to population stability or growth, but reduction to < 2 wolves/1000 km² is required to elicit the strongest possible population response (Serrouya et al. 2019). The results achieved in the South Peace treatment herds have equalled, or surpassed, those achieved by other

jurisdictions that have implemented wolf reduction to enhance caribou populations (Farnell and McDonald 1988, Boertje et al. 1996, Bergerud and Elliot 1998, Hayes et al. 2003, Hervieux et al. 2014). Wolf reduction as a caribou recovery tool may be particularly valuable for herds that require immediate recovery actions while the ultimate causes of caribou population declines (e.g. habitat alteration) are addressed.

Caribou calf recruitment remained lower than expected in the Klinse-Za herd despite intensive wolf reduction. This was further evidenced when comparing the survival rates of calves released from the maternal pen to those of the unpenned calves in the population (McNay et al. 2019). For example, in 2018, the annual survival rate of penned calves following their release was 77% versus only 17% survival amongst unpenned calves. Calf survival had increased substantially in the other South Peace treatment herds following effective wolf reduction, which suggests there were factors specific to the Klinse-Za range that were inhibiting calf survival. Bears (*Ursus spp.*) are believed to be abundant throughout the Klinse-Za range and have been shown to be significant predators of calf caribou (Adams et al. 1995). Similarly, calf recruitment observed in the South Narraway herd following one year of wolf reduction was lower than expected; however, further years of treatment may be required in order to obtain stronger inferences. It would be beneficial to implement research studies to investigate the cause of low survival of unpenned wild calves in the Klinse-Za caribou herd.

Prior to population declines, it is believed that approximately 1,000 caribou inhabited the South Peace caribou range (approximately 800 amongst the Klinse-Za, Kennedy Siding, and Quintette herds, and 200 within the South Narraway herd; Ministry of Environment 2013, Seip and Jones 2014). Assuming continued annual population growth of approximately 15% (achieved solely through wolf reduction or in conjunction with other management initiatives), the treatment herds could double in size by 2026-2027 and the objective of 800 caribou in the Pine and Quintette LPUs could be met by 2036-2037. In order for those caribou herds to achieve a self-sustaining status, however, the ultimate causes of declines (i.e. industrial landscape change and apparent competition) must be addressed through habitat protection, recovery, and restoration. It is too early to determine whether the population target of 200 caribou in the South Narraway range can be achieved through wolf reduction.

Annual Wolf Recovery

Previous wolf reduction programs have experienced challenges with rapid population growth or recolonization by wolves on an annual basis (Bergerud and Elliot 1998, Hayes et al. 2003, Hervieux et al. 2014). During years of intensive wolf reduction in the South Peace, there were few or no breeding pairs remaining in the treatment area following the winter removal efforts (as determined by closely documenting wolf presence throughout the aerial reduction efforts), which reduced the likelihood of litter production during the spring. Thus, the subsequent recovery of wolves was occurring primarily through colonization of new wolves into the treatment area. Wolf recolonization is presumed to occur rapidly when primary prey within the treatment area remains abundant, and where there are few geographic barriers inhibiting wolf movement into the treatment area. In the South Peace, the low presence of linear disturbance (relative to many Boreal caribou ranges) combined with the mountainous landscape bordered by large waterbodies (on the western and northern extent) may slow the rate of wolf recolonization relative to those rates observed in highly disturbed and less mountainous landscapes (Hervieux et al. 2014). Additionally, wolf reduction occurring to the east of the treatment area in Alberta may also reduce the rate of wolf dispersal from neighbouring jurisdictions.

Wolf reductions in the South Peace have shown that annual population growth and recolonization rates by wolves were significant when less than 80% of wolves were removed during the winter. Following winters of intensive reduction (greater than 90% removal), the subsequent wolf populations had only amounted to approximately 65% of the previous year's population. Prior to reduction efforts in the winter of 2018-2019, the

wolf population within the treatment area was only 38% of the initial wolf population present prior to the program's commencement in 2014-2015. These results suggest that continued, intensive wolf reduction can decrease the overall presence of wolves in a treatment area on an annual basis. As reported in similar studies (i.e. Bergerud and Elliot 1998), over the course of multiple years of reduction efforts wolves were recolonizing in smaller pack sizes; an expected result based on the breeding and dispersing behaviour of wolves (Fuller et al. 2003). Small and numerous wolf packs increase the difficulty of removing a high proportion of wolves, as they are more challenging to locate than larger packs and have less-established territories.

Primary Prey

Relative to caribou recovery, an increase in primary prey populations would be considered an undesirable side effect of wolf reductions, due to the apparent competition hypothesis (Seip 1992). In other jurisdictions, wolf reduction has been used as a management tool to deliberately increase primary prey populations for species such as moose, elk (*Cervus canadensis*), and thimhorn sheep (*Ovis dalli*) and has been successful in doing so in many cases (Boertje et al. 1996, Bergerud and Elliot 1998, Hayes et al. 2003, Keech et al. 2011). Based on the results of those programs, it is expected that primary prey species in the South Peace, particularly moose and to a lesser extent deer (*Odocoileus spp.*) and elk, would respond at an equal or greater level to that of the caribou populations. A lack of baseline data for moose populations across the treatment area makes for difficult interpretations of the response to wolf reduction; however, the available data do not suggest as strong of a response as expected. Although survival rates of adult moose have likely increased and positive population growth has occurred (i.e. within the Klinse-Za range; Sittler 2019), there has been a lesser response in calf moose survival and recruitment. This may indicate that there are other factors slowing the rate of population growth amongst moose, such as bear predation on neonate calves, or health factors (Kuzyk et al. 2018). Other jurisdictions within BC have recently reported high moose population growth and calf recruitment in response to wolf reduction (Serrouya and Legebokow 2018).

The response of primary prey to wolf reduction requires further monitoring, and wildlife managers must consider options for managing primary prey in order to reduce the recovery rates by wolves. This may be achieved through liberalized licensed hunting opportunities or managing habitat such that it is less suitable for primary prey species. Messier's (1994) numeric response model predicts that moose densities below 0.13 moose/km² are necessary for wolves to meet the threshold associated with caribou population stability (6.5 wolves/1000 km²; Bergerud and Elliot 1986). Similar prey biomass equations estimate that moose densities below 0.3 moose/km² are required for wolf densities below 6.5 wolves/1000 km² (Fuller 1989, Wilson 2009), which is comparable to Bergerud's (1996) research which suggested caribou cannot persist when moose densities exceed 0.2–0.3 moose/km². Furthermore, moose densities of less than 0.2 moose/km² have been associated with reduced wolf recruitment (Messier 1985, Serrouya et al. 2017). Moose densities of greater than 0.2–0.3 moose/km² in the South Peace may not impede caribou recovery efforts so long as intensive wolf reduction continues to be used as a management option. However, it would be detrimental to allow moose abundance to occur beyond such densities, as it would continue to facilitate annual wolf recovery, and ultimately hinder caribou populations from achieving self-sustaining status. Furthermore, wolf reduction is not viewed as a long-term management tool, thus moose abundance should be addressed through active moose population management and caribou habitat recovery. It must be recognized that there are significant social and logistical challenges in maintaining or reducing moose densities through licensed hunting, particularly due to First Nations' desire for abundant moose populations to meet their food, social, and ceremonial rights.

Links to Other Caribou Recovery Initiatives

Wolf reduction as a caribou recovery tool is likely to achieve the greatest results when it occurs in conjunction with other recovery initiatives that address underlying causes of caribou population declines (Serrouya et al. 2019). The long-term recovery of caribou populations requires the recovery of caribou habitat to a state that resembles pre-disturbance conditions. Habitat change that has benefited primary prey species or enhanced wolf movement across the landscape must be addressed in order for the long-term caribou recovery objectives to be met. Across the South Peace caribou range, habitat protection and restoration initiatives are underway and continue to be pursued (MFLNRO 2017, Woods and McNay 2019). Such initiatives are necessary to support caribou recovery, particularly if and when wolf reduction is halted as a management tool for South Peace caribou. Furthermore, social acceptance for wolf reduction is contingent on the Ministry's ability to demonstrate commitment to addressing the ultimate causes of caribou population declines. Wolf reduction should be viewed as a short-term recovery tool that supports South Peace caribou herds while habitat conditions improve.

There are other short-term caribou recovery measures occurring in conjunction with wolf reduction across the South Peace caribou ranges. In the Kennedy Siding herd, supplemental feeding of caribou during the autumn and early winter has occurred for several years (Heard and Zimmermann 2018). While the positive effects of feeding caribou are somewhat inferred (i.e. improved nutritional status), the Kennedy Siding herd did not show a measurable population response to supplemental feeding in the absence of wolf reduction. Within the Klinse-Za caribou range, an ongoing maternal penning program has been underway since 2014 (McNay et al. 2019). Initial results from the maternal penning efforts, prior to the initiation of wolf reduction, suggested that penning was ineffective if wolf populations were not reduced. During the program's first year, five of nine calves were killed soon after their release from the pen (a higher mortality rate than unpenned calves) and adult mortality remained high (Seip and Jones 2016). However, when combined with intensive wolf reduction in subsequent years, calf survival has increased (Seip and Jones 2018, McNay et al. 2019). Calf survival of penned calves has been higher than that of unpenned calves in all years with concurrent wolf reduction, suggesting that the maternal penning efforts are contributing additional calves to the population that may otherwise have perished. The maternal penning project does retain some risk, however. The repeated capture, retention, and rearing of caribou can be stressful, result in injury and death, and may have short- and long-term health and behavioural implications. Additionally, the costs associated with maternal penning relative to population growth are significantly higher when compared to wolf reduction alone (i.e. the cost per caribou added to the population through maternal penning is approximately one order of magnitude greater than that of wolf reduction). In the South Peace, maternal penning is not supported, nor would it be expanded to include new caribou herds, in the absence of intensive, concurrent wolf reduction. Furthermore, the Kennedy Siding and Quintette caribou herds have shown comparable or greater annual population growth and calf recruitment than the Klinse-Za herd through wolf reduction alone.

Operational Considerations

Overall, the delivery of the wolf reduction program in the South Peace has increased in efficacy and efficiency over time as crews have gained more experience and familiarity with the treatment area and as the operational oversight by Ministry staff has increased. Cost efficiency is dependent on the experience and proficiency levels of the removal crews, weather conditions, and the abundance and pack sizes of wolves within the treatment area. The effort required to reduce wolf densities to a low level is substantial. Generally, greater than 225 hours of helicopter flight time are flown to achieve successful wolf reduction across the South Peace treatment area. This necessitates a substantial financial commitment, as well as commitment of time and capacity from the aerial removal crews and Ministry staff. A sufficient budget must be forecasted each year, as well as a multi-year funding commitment. Ministry staff capacity must also be forecasted, and wolf reduction crews with

demonstrated proficiency must be identified and procured. The results reported during this program suggest that wolf reduction must be very intensive, and to implement a program that is anything less than intensive would be considered unethical. Scientific rigour is required to deliver the removal aspects of this program and accurately measure the response of both wolves and caribou to wolf reduction. Wolf reduction has proved to be most efficient and effective when conducted under optimal weather conditions that facilitate the tracking, locating, capturing and radio collaring, and ultimately the removal of the majority of wolves within the treatment area. The intensity of wolf reductions achieved during this program has generally exceeded those reported in other jurisdictions where wolf reduction has occurred (Bergerud and Elliot 1998, Hayes et al. 2003, Hervieux et al. 2014).

A critical factor to a successful program is the radio collaring of most, if not all, wolf packs within the treatment area. The deployment of radio-collars greatly reduces search times when locating wolf packs, and aids in the facilitation of complete pack removal. Wolves captured and fitted with new, active radio-collars were generally left alive following the reduction efforts in order to facilitate relocation of packs the following winter. Under most circumstances, the lone radio-collared wolves had not bred successfully, thus were not part of new family units the following winter. Furthermore, it was rare that these lone wolves were accepted into new packs that may have colonized the treatment area. The most common scenario was that radio-collared wolves partnered with other individuals or pairs of wolves to form new pairs or small groups. Most years, approximately half of the radio-collared wolves survived or remained within the treatment area by the following winter. Although the removal of the radio-collared wolves annually would have resulted in a lower wolf density at the end of each winter's reduction efforts, the value of leaving those wolves is believed to outweigh the benefits of removing them.

The concurrent ground removal programs implemented by First Nations (in the Klinse-Za range) provided an additional source of wolf reduction, as well as an opportunity to collaborate with local First Nations communities on caribou recovery initiatives. In treatment areas where there were no ground removal programs (i.e. Kennedy Siding and Quintette), the aerial removal on its own was shown to be sufficiently effective. There were also risks of actively trapping in conjunction with aerial removal, the primary risk being that a radio-collared individual could have been accidentally trapped and killed. Due to the elusive nature of wolves, their large home ranges and propensity for remote, inaccessible locations, trapping and hunting is unlikely to achieve wolf reduction targets as a standalone measure. Furthermore, the reduction of wolves through hunting or trapping has been shown to have little effect on wolf populations (Webb et al. 2011). Given these factors, it is unlikely that wolf reduction through hunting and/or trapping is sufficient to elicit positive responses in caribou populations.

The cost effectiveness of the South Peace wolf reduction can be measured using several variables. Most notably, the cost per caribou added⁵ to the population following three years of effective wolf reduction equates to approximately \$11,000 per caribou. During that time, there has been one caribou added to the population for every 2.9 wolves that have been removed. The cost per wolf removed (through aerial shooting) has averaged approximately \$5,100 over the past five years; this cost has fluctuated annually and is dependent on the proficiency of the removal crews, weather conditions, abundance of wolves, and pack sizes (where more effort is required over time to remove numerous, but small packs). The overall cost to deliver wolf reduction in the South Peace treatment area may be lessened over time, provided the wolf recovery rate remains low, as was observed in 2018-2019. The program costs and number of wolves removed are largely independent of the

⁵ Calculated by identifying the number of caribou added to the population as well as the number of caribou that would have been lost from the population in the absence of wolf reduction

caribou population size; thus, the cost and number of wolves removed annually may remain constant over time, but the number of caribou added annually will increase as the caribou population size increases (assuming similar lambda results are achieved). If primary prey populations (and their preferred habitats) are managed and caribou populations continue to increase at their current rate, it is possible that a lesser effort to reduce wolves could be applied on an annual basis, or wolf reductions could occur on two- or three-year cycles. As the caribou population increases, the cost per caribou added and the number of caribou added per wolf removed (presuming similar population growth rates are achieved)

During Years 1 and 2 of the reduction program, there were attempts made to retrieve the majority of wolf carcasses by helicopter in order to provide pelts to local First Nations. It soon became evident that the retrieval of carcasses was highly inefficient (required much additional helicopter flight time and time on the ground by removal crews), increased the safety risks for removal crews, and had low uptake from First Nations relative to utilization of the pelts. In the subsequent years of the program, Ministry staff and removal crews have coordinated the ground retrieval of wolf carcasses that were relatively accessible, which were then distributed to local First Nations. This has been a much more efficient method of retrieving and distributing carcasses, and has generally provided a sufficient number of pelts to First Nations to support their social and ceremonial interests.

Aerial shooting is the most effective method of reducing wolf densities over large geographic areas while eliminating the risk of bycatch and ensuring the highest likelihood of quickly dispatching wolves. The humaneness of wolf removal in the South Peace has been examined more thoroughly as Ministry staff has become more involved in the operational delivery. In 2018-2019, humaneness was examined by Ministry staff by documenting shooting proficiency, shot locations, and subsequent dispatch times of a large subsample of wolves removed during program delivery (including the South Peace and two other treatment areas in the region). Of the documented subsample of 98 wolf removals, the vast majority of wolves were dispatched instantaneously or within seconds following one well-placed shot or a quick succession of multiple shots. Only six wolves took longer than 30-seconds to expire after an initial shot, and only one wolf was never visually confirmed to have expired after being shot (although it appeared to have expired out of sight in a tree well). It is important that Ministry staff, working closely with the removal crews, continue to document the shooting proficiency and effectiveness during wolf reduction programs and adjust methods as necessary to ensure wolf reduction occurs at the highest possible level of humaneness.

5. Conclusion and Recommendations

The South Peace aerial wolf reduction program has demonstrated conclusively that the reduction of wolves across the treatment area has had a strong, positive effect on the caribou populations. Wolf reduction is a management tool that must be used responsibly and ethically, and implemented with the highest standards for humaneness and scientific rigour. Reducing wolf populations may be the most effective interim management measure for halting and reversing caribou population declines over the short-term while the ultimate causes of such declines are addressed through habitat protection and restoration and primary prey management. Wildlife managers and Ministry decision-makers should consider the following recommendations that have been identified during the five-year review of wolf reduction in the South Peace region:

1. Continue the intensive reduction of wolves across the South Peace treatment areas until caribou populations approach a self-sustaining status (approximately 1,000 individuals)
 - Consider approval of an additional five years of wolf reduction, followed by a comprehensive review of the program

- Implement multiple management tools, including habitat protection and restoration, concurrently to help address the ultimate causes of population declines
- 2. Consider lessening the wolf reduction intensity if wolf recovery remains low and caribou continue to trend towards self-sustaining levels
 - Contingent on habitat restoration, protection, and maintenance of primary prey populations
- 3. Consider a “maintenance” approach to wolf reductions for the South Narraway caribou herd, provided wolf recovery remains low
 - This may be achieved through reducing the annual reduction effort, or conducting reductions on a two- to three-year cycle
- 4. Continue to use experienced and proficient removal crews that can be trusted to deliver wolf reduction in a professional and humane manner
- 5. Maintain Ministry staffs’ responsibility and role in project coordination, operational oversight, and field involvement
- 6. Continue the intensive monitoring of caribou populations in response to wolf reductions, including the documentation of annual population estimates, population trend, calf recruitment, and adult female survival rates
- 7. Monitor the response by primary prey to wolf reductions, establish target densities for primary prey species (i.e. moose) within the treatment areas, and implement measures that can be used to achieve and maintain those objectives
 - Consider using licensed hunting to manage for moose population densities between 0.2–0.3 moose/km²
- 8. Consider habitat protection and restoration measures such that habitat gain exceeds habitat loss within and adjacent to caribou core ranges
- 9. Continue to rigorously document the humaneness of the wolf reduction efforts
- 10. Continue to support the ground retrieval of carcasses to be distributed to local First Nations
- 11. Continue to implement a collaborative approach to wolf reduction by supporting ground trapping programs by local First Nations
- 12. Consider research opportunities to investigate causes of caribou calf mortality (i.e. predation by bears or other predators, health-related causes, etc.)
- 13. Continue to implement the Ministry’s safe work practices for aerial shooting of wolves and net-gunning capture of wolves

6. Literature Cited

- Adams, L. G., F. J. Singer, and B. W. Dale. 1995. Caribou calf mortality in Denali National Park, Alaska. *Journal of Wildlife Management* 59:584-594.
- Bergerud, A.T. 1992. Rareness as an antipredator strategy to reduce predation risk for moose and caribou. *In* *Wildlife 2001: populations*. Edited by D.R. McCullough and R.B. Barrett. Elsevier, London, UK. pp 1008–1021.
- Bergerud, A.T., and J.P. Elliot. 1986. Dynamics of caribou and wolves in northern British Columbia. *Canadian Journal of Zoology* 64:1515–1529.
- Boertje, R. D., P. Valkenburg, and M. E. McNay. 1996. Increases in moose, caribou, and wolves following wolf control in Alaska. *The Journal of Wildlife Management* 60: 474–489.

- COSEWIC. 2011. Designatable units for caribou (*Rangifer tarandus*) in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa, ON, Canada. 88 pp.
- Environment Canada. 2014. Recovery strategy for the woodland caribou, southern mountain population (*Rangifer tarandus caribou*) in Canada. Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa, ON, Canada. 68 pp.
- Farnell, R., and J. McDonald. 1988. The demography of Yukon's Finlayson caribou herd, 1982 to 1987. Progress report, Yukon Renewable Resources, Whitehorse, YK, Canada.
- Fuller, T. K. 1989. Population dynamics of wolves in north-central Minnesota. *Wildlife Monographs* 105:3–41.
- Fuller, T. K., L. D. Mech, and J. F. Cochrane. 2003. Wolf population dynamics. *In* *Wolves: behavior, ecology, and conservation*. Edited by D. L. Mech and L. Boitani. University of Chicago Press, Chicago IL, USA. pp 161-19.1
- Gustine, D.D., K.L. Parker, R.J. Lay, M.P. Gillingham, D.C. Heard. 2006. Calf survival of woodland caribou in a multi-predator ecosystem. *Wildlife Monograph* 165:1–32.
- Hatter, I.W., and W. A. Bergerud. 1991. Moose recruitment, adult mortality and rate of change. *Alces*, 27:65–73.
- Hayes, R.D., R. Farnell, R.M. Ward, J. Carey, M. Dehn, G.W. Kuzyk, A.M. Baer, C.L. Gardner, and M. O'Donoghue. 2003. Experimental reduction of wolves in the Yukon: ungulate responses and management implications. *Wildlife Monographs* 152:1–35.
- Heard, D., and K. Zimmerman. 2018. Supplemental feeding of Kennedy Siding caribou, September 2017 to January 2018. Report for the Peace Northern Caribou Program. 21p.
- Hervieux, D., M. Hebblewhite, D. Stepinsky, M. Bacon, and S. Boutin. 2014. Managing wolves (*Canis lupus*) to recover woodland caribou (*Rangifer tarandus caribou*) in Alberta. *Canadian Journal of Zoology* 92:1029–1037.
- Keech, M. A., M. S. Lindberg, R. D. Boertje, P. Valkenburg, B. D. Taras, T. A. Boudreau, and K. B. Beckmen. 2011. Effects of predator treatments, individual traits, and environment on moose survival in Alaska. *Journal of Wildlife Management* 75:1361–1380.
- Klaczek, M., S. Marshall, D. Heard, and A. Batho. 2016. Density and composition of moose (*Alces alces*) within the Pine and Mackenzie study area (WMU 7-23A and 7-30) in north-central British Columbia: winter 2015-2016. BC Ministry of Forests, Lands and Natural Resource Operations, Prince George, BC, Canada.
- Kuzyk, G. 2002. Wolf distribution and movements on caribou range in West-central Alberta. M.Sc. thesis, University of Alberta.
- Kuzyk, G., S. Marshall, C. Procter, H. Schindler, H. Schwantje, M. Gillingham, D. Hodder, S. White, and M. Mumma. 2018. Determining factors affecting moose population change in British Columbia: testing the landscape change hypothesis. 2018 Progress Report: February 2012–April 2018. BC Ministry of Forests, Lands, Natural Resource Operations and Rural Development. Victoria, BC, Canada, Working Report No. WR-126. Pp. 64.

- McNay, R.S., M. Erickson, and L. Giguere. 2019. Enhancing calf survival to help avert extirpation of the Klinse-Za caribou herd: annual report. Wildlife Infometrics Inc. Report No. 671. Wildlife Infometrics Inc., Mackenzie, BC, Canada.
- Messier, F. 1985. Social organization, spatial distribution, and population density of wolves in relation to moose density. *Canadian Journal of Zoology* 63:1068–1077.
- Messier, F. 1994. Ungulate population models with predation: a case study with the North American moose. *Ecology* 75:478–488.
- Ministry of Environment. 2013. Implementation plan for ongoing management of South Peace northern caribou (*Rangifer tarandus caribou* pop. 15) in British Columbia. British Columbia Ministry of Environment technical report, Victoria, BC, Canada.
- Ministry of Forests, Lands and Natural Resource Operations. 2017. British Columbia's Quintette strategic action plan – southern mountain caribou herd recovery. British Columbia Ministry of Forests, Lands and Natural Resource Operations technical report, Victoria, BC, Canada.
- Seip, D. R. 1992. Factors limiting woodland caribou populations and their relationships with wolves and moose in southeastern British Columbia. *Canadian Journal of Zoology* 70:1494–1503.
- Seip, D.R., and E. Jones. 2014. Population status of caribou herds in the Central Mountain Designatable Unit within British Columbia, 2014. Unpublished report. BC Ministry of Environment.
- Seip, D.R., and E. Jones. 2016. Population status of central mountain caribou herds within British Columbia, 2015. Unpublished report. BC Ministry of Environment.
- Serrouya, R.B., and C. Legebokow. 2018. Moose population monitoring in the Lake Revelstoke Valley (Management Units 4-38 and 4-39), February 2018. Unpublished Report, Columbia Mountains Caribou Project and BC Ministry of Forests, Lands, and Natural Resource Operations.
- Serrouya, R., B. N. McLellan, H. van Oort, G. Mowat, and S. Boutin. 2017. Experimental moose reduction lowers wolf density and stops decline of endangered caribou. *PeerJ* 5:e3736.
- Serrouya R., D.R. Seip, D. Hervieux, B.N. McLellan, R.S. McNay, R. Steenweg, D.C. Heard, M. Hebblewhite, M. Gillingham, and S. Boutin. 2019. Saving endangered species using adaptive management. *PNAS* 116: 6181–6186.
- Sittler, K.L. 2019. Moose Limiting Factors Investigation: Annual Report 2018-19. Wildlife Infometrics Inc. Report No. 678. Wildlife Infometrics Inc., Mackenzie, BC, Canada.
- Sittler, K.L. and R.S. McNay. 2018. Moose (*Alces americanus*) population density and composition around the Clearwater and Moberly River drainages (WMU 7-31), Northeastern BC. Wildlife Infometrics Inc. Report No. 623. Wildlife Infometrics Inc., Mackenzie, BC, Canada.
- Webb, N. F., J. R. Allen, and E. H. Merrill. 2011. Demography of a harvested population of wolves (*Canis lupus*) in west-central Alberta, Canada. *Canadian Journal of Zoology* 89:744–752.

- Wilson, S. F. 2009. Recommendations for predator-prey management to benefit the recovery of mountain caribou in British Columbia. Ministry of Environment – Environmental Stewardship Division. Victoria, BC, Canada.
- Woods A.D., and R.S. McNay. 2019. Klinse-Za/Scott East caribou habitat restoration: 2018-19 annual report. Wildlife Infometrics Inc. Report No. 669. Wildlife Infometrics Inc., Mackenzie, BC, Canada.

From: [Hansen, Inge-Jean FLNR:EX](#)
To: [Bridger, Michael C FLNR:EX](#)
Subject: RE: South Peace Caribou Recovery - Experimental Wolf Reduction report
Date: Wednesday, September 25, 2019 12:35:43 PM
Attachments: [image001.png](#)
[image003.png](#)

Great work Mike! I haven't read through it all yet, but on first scan I would say you have done a great job.

Appreciate the rigorous review you sought out.

I-J

From: Bridger, Michael C FLNR:EX
Sent: Wednesday, September 25, 2019 11:46 AM
To: Vince, Karrilyn M FLNR:EX; Chisholm, Diane FLNR:EX; Jones, Aviva FLNR:EX; Hansen, Inge-Jean FLNR:EX; Gagne-Delorme, Audrey FLNR:EX; Harvey, Kerry FLNR:EX; Watt, Kevin FLNR:EX; Lavallee, Michel FLNR:EX; Peck, Kristen ENV:EX; Nicholls, Emony FLNR:EX; Sinitsin, Chelsea FLNR:EX; Kabzems, Richard FLNR:EX
Cc: Pelletier, Agnes FLNR:EX; Schilds, Scott FLNR:EX; Scheck, Joelle L FLNR:EX
Subject: South Peace Caribou Recovery - Experimental Wolf Reduction report

Hey everyone,

Just wanted to pass this along to the RM team. I've attached the 5-year summary report for wolf reduction in the South Peace to support caribou recovery. This report has undergone extensive reviews from nine biologists both internal and external to government. As it stands, this is the best current science we have in BC demonstrating the response by caribou to intensive wolf reduction, and are arguably the strongest results that have been achieved for caribou recovery in North America through the use of predator reduction. This report has been approved for external distribution, so it may be a valuable resource to support conversations with First Nations and stakeholders. I will save a copy on the I:drive in the wildlife reports folder as well.

And as an FYI, the long-awaited NE predator management plan was approved by the North Area ADM recently, following approval from the North Area REDs last February. We'll be collecting the necessary signatures in the next week or two, and then I can distribute that document as well.

Let me know if you have any comments or questions.

Cheers,

Mike

Mike Bridger, M.Sc., R.P.Bio.



Wildlife Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development
Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7
Ph: 250-787-3294



Page 80 of
92
92-AR-
2019-96488

Page 81 of
92 FNR-
2019-96488

From: Bridger, Michael C FLNR:EX
To: Peel, Darcy E FLNR:EX; Scheck, Joelle L FLNR:EX; DeGroot, Leo FLNR:EX; Wiebe, Heather I FLNR:EX; Sobieszczyk, Beth FLNR:EX; Muter, David FLNR:EX; Hughes, Janet FLNR:EX; Vince, Karrilyn M FLNR:EX; Chisholm, Diane FLNR:EX; Davis, Celine ENV:EX; Zimmerman, Kathi ENV:EX; Huck, Michael FLNR:EX; Pelletier, Agnes FLNR:EX; Schilds, Scott FLNR:EX; Watt, Kevin FLNR:EX; Friedrichsmeier, Christine FLNR:EX; Ethier, Tom FLNR:EX; Haslam, David GCPE:EX; Makarowski, Dawn GCPE:EX; Laroche, Russ FLNR:EX; Ernst, Bevan FLNR:EX; Reid, Aaron FLNR:EX; Shores, Carolyn FLNR:EX; White, Shane FLNR:EX; Hoffos, Robin FLNR:EX; Skerik, David FLNR:EX; Jacklin, James FLNR:EX
Subject: RE: Energetic city Wolf cull article
Date: Thursday, October 10, 2019 10:26:51 AM
Attachments: [image001.png](#)
[image003.png](#)
[image004.png](#)

And a few others, mainly positive. Although the habitat piece comes in to question particularly in the Vancouver Sun article.

<https://vancouversun.com/news/local-news/caribou-numbers-up-49-per-cent-after-four-year-wolf-cull>

<https://www.abbynews.com/news/wolf-kill-working-in-b-c-caribou-recovery-ministry-study-shows/>

Mike Bridger
Wildlife Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Peel, Darcy E FLNR:EX
Sent: Wednesday, October 9, 2019 9:56 PM
To: Scheck, Joelle L FLNR:EX; DeGroot, Leo FLNR:EX; Wiebe, Heather I FLNR:EX; Sobieszczyk, Beth FLNR:EX; Muter, David FLNR:EX; Hughes, Janet FLNR:EX; Vince, Karrilyn M FLNR:EX; Chisholm, Diane FLNR:EX; Davis, Celine ENV:EX; Zimmerman, Kathi ENV:EX; Bridger, Michael C FLNR:EX; Huck, Michael FLNR:EX; Pelletier, Agnes FLNR:EX; Schilds, Scott FLNR:EX; Watt, Kevin FLNR:EX; Friedrichsmeier, Christine FLNR:EX; Ethier, Tom FLNR:EX; Haslam, David GCPE:EX; Makarowski, Dawn GCPE:EX; Laroche, Russ FLNR:EX; Ernst, Bevan FLNR:EX; Reid, Aaron FLNR:EX; Shores, Carolyn FLNR:EX; White, Shane FLNR:EX; Hoffos, Robin FLNR:EX; Skerik, David FLNR:EX; Jacklin, James FLNR:EX
Subject: Re: Energetic city Wolf cull article

Thank you joelle, nice to get some positive news out.

Darcy

Sent from my Samsung Galaxy smartphone.

----- Original message -----

From: "Scheck, Joelle L FLNR:EX" <Joelle.Scheck@gov.bc.ca>
Date: 2019-10-09 5:36 p.m. (GMT-08:00)
To: "Peel, Darcy E FLNR:EX" <Darcy.Peel@gov.bc.ca>, "DeGroot, Leo FLNR:EX" <Leo.DeGroot@gov.bc.ca>, "Wiebe, Heather I FLNR:EX" <Heather.Wiebe@gov.bc.ca>, "Sobieszczyk, Beth FLNR:EX" <Beth.Sobieszczyk@gov.bc.ca>, "Muter, David FLNR:EX" <David.Muter@gov.bc.ca>, "Hughes, Janet FLNR:EX" <Janet.Hughes@gov.bc.ca>, "Vince, Karrilyn M FLNR:EX" <Karrilyn.Vince@gov.bc.ca>, "Chisholm, Diane FLNR:EX" <Diane.Chisholm@gov.bc.ca>, "Davis, Celine ENV:EX" <Celine.Davis@gov.bc.ca>, "Zimmerman, Kathi ENV:EX" <Kathi.Zimmerman@gov.bc.ca>, "Bridger, Michael C FLNR:EX" <Michael.Bridger@gov.bc.ca>, "Huck, Michael FLNR:EX" <Michael.Huck@gov.bc.ca>, "Pelletier, Agnes FLNR:EX" <Agnes.Pelletier@gov.bc.ca>, "Schilds, Scott FLNR:EX" <Scott.Schilds@gov.bc.ca>, "Watt, Kevin FLNR:EX" <Kevin.Watt@gov.bc.ca>

Subject: Fwd: Energetic city Wolf cull article

Fyi

Sent from my iPhone

Begin forwarded message:

From: "Cote, Kelli FLNR:EX" <Kelli.Cote@gov.bc.ca>
Date: October 9, 2019 at 4:36:37 PM MST
To: "Scheck, Joelle L FLNR:EX" <Joelle.Scheck@gov.bc.ca>
Subject: Energetic city Wolf cull article

FYI

<https://www.energeticcity.ca/2019/10/caribou-population-up-by-almost-50-percent-following-five-year-wolf-cull-in-south-peace/>



Kelli Côté, R.P.Bio.

Senior Caribou Project Manager
BC CARIBOU RECOVERY PROGRAM - NE Team
Ministry of FLNRORD – NE Region

Suite 300, 10003 – 110th Avenue | Fort St John, BC V1J 6M7
Phone: 250-787-3479



Page 84 of
92
92-AR-
2019-96488

Page 85 of 92
FNR-2019-
96488

To: [XT:Serrouya, Rob FLNR:IN](#)
Subject: RE: 5-Year Wolf Reduction Report
Date: Wednesday, November 6, 2019 11:09:21 AM
Attachments: [image001.png](#)
[image004.png](#)

From: Robert Serrouya [mailto:serrouya@ualberta.ca]
Sent: Tuesday, August 6, 2019 3:11 PM
To: Bridger, Michael C FLNR:EX
Subject: Re: 5-Year Wolf Reduction Report

Hiya - could you please send me the shapefile polygon, or any GIS type format, of your red areas in fig 2? Just needed for the moose / caribou extension note I'm working on with Burwash & Kuzyk. Thanks

On Tue, Aug 6, 2019 at 2:07 PM Bridger, Michael C FLNR:EX
<Michael.Bridger@gov.bc.ca> wrote:

This is great Rob, thanks!

Just went through and made some of the simple changes and edits, now I'm just going through again and address some of your bigger-picture thoughts. Definitely some strong considerations given, and I will try to incorporate most them.

The humane piece is a tricky one, agreed. We did try to assess this more thoroughly this past year.

You should bill us for your time. I have money for it. Can you send me an invoice?

Mike Bridger
Caribou Recovery Biologist
FLNRORD Northeast Region
Phone: (250)787-3294



From: Robert Serrouya [mailto:serrouya@ualberta.ca]
Sent: Tuesday, August 6, 2019 11:54 AM
To: Bridger, Michael C FLNR:EX
Subject: Re: 5-Year Wolf Reduction Report

Hi Mike

Here is my VERY brief read of your report. It is an impressive piece of work. Very clear, great data. I wish I could spend more time on it this month. I'm happy to help in any capacity in the future. Call if needed.

Thanks

Rob

On Fri, Aug 2, 2019 at 9:47 AM Bridger, Michael C FLNR:EX

<Michael.Bridger@gov.bc.ca> wrote:

Latest version attached.

Mike Bridger

Caribou Recovery Biologist

FLNRORD Northeast Region

Phone: (250)787-3294



From: Robert Serrouya [mailto:serrouya@ualberta.ca]

Sent: Tuesday, July 30, 2019 8:21 AM

To: Bridger, Michael C FLNR:EX

Subject: Re: 5-Year Wolf Reduction Report

You bet, I'll have a quick look . If you have a sec to call me, that would be great. thanks



On Mon, Jul 29, 2019 at 1:30 PM Bridger, Michael C FLNR:EX

<Michael.Bridger@gov.bc.ca> wrote:

No pressure, but do you think you'll have a chance to read thru the report this week?

I just need to update folks with the timeline.

Mike Bridger

Caribou Recovery Biologist

FLNRORD Northeast Region

Phone: (250)787-3294



From: Bridger, Michael C FLNR:EX

Sent: Thursday, July 18, 2019 2:59 PM

To: XT:Serrouya, Rob FLNR:IN

Subject: RE: 5-Year Wolf Reduction Report

Totally understand about the timeline. The whole thing is a bit rushed.

Would love to get your eyes on it still, even just at a high-level to hear your thoughts on the recommendation section, etc. If you have time and want to take a look through, that would be great. I've attached the draft here. If you spend an hour or two on it, I'd still want to compensate you for your time, so just send me an invoice with an hourly rate if that works for you?

I'll ask again about the timeline, but I'm away for nearly all of September. That would push things back to October, and that might be too late for the decision makers. They want to use

this finalized report to guide their decisions around wolf control expansion into the newly proposed areas, hence the rush.

Mike Bridger

Caribou Recovery Biologist

FLNRORD Northeast Region

Phone: (250)787-3294



From: Robert Serrouya [mailto:serrouya@ualberta.ca]

Sent: Thursday, July 18, 2019 6:53 AM

To: Bridger, Michael C FLNR:EX

Subject: Re: 5-Year Wolf Reduction Report

Hey Mike

This sounds very interesting. I like the idea of an internal document followed by something more general. This would also likely fit within my workplan for work I am doing for Michael Burwash. The main hang up at this point is timeline - I can't commit to a 30-page review by mid August (learning to manage time has been a major challenge for me!). I am writing up the 5 year penning report which is due sept 15 and is very involved. The best I could do is a 1 hr skim but not sure that would be of much help. Would 1 more month be possible? And yes would be happy to help, if I can contribute anything new, to the eventual publication.

Talk soon

Rob

On Tue, Jul 16, 2019 at 2:29 PM Bridger, Michael C FLNR:EX
<Michael.Bridger@gov.bc.ca> wrote:

Hey Rob,

I've written a draft of the 5-Year Experimental Wolf Reduction for Central Mountain Caribou. It has been reviewed internally by a few folks, including some Caribou Science Team members. We're looking to have one or two folks provide an external review as well. I'm thinking this report would be of interest to you.

As it stands now, it's somewhat written as an internal-facing document, just because it does include some details around contractor proficiency, budgets, etc. After all of the final reviews, I'll probably create a version that is outward facing. The caribou program would like to be able to share the document broadly by late-August I believe.

So, would you be interested in reviewing it? It's about 30 pages, and I would send you the data as well. My timeline for receiving edits and comments would be around the 1st or 2nd week of August, because I'm taking off for a while starting the second

half of August. We would just do a direct invoice with you, if you're able to provide an hourly or daily rate for your time.

One thing to keep in mind if you review it is the potential to turn this work into a scientific publication. I would like to get the 5th year of caribou response data first, just to increase the sample size. But would be curious to get your opinion if you think there would be enough good content for a publication. You come to mind as a potential co-author if we do decide to go down that road. In the current report, the data and results may not be displayed at a level suitable for academic purposes (which I think is okay for this version of the report, it's supposed to be somewhat 'easy-reading'), but it would be good to at least think of some ways to strengthen the reporting of results. For example, I haven't come up with a way to portray ranges of error around some of the estimates.

Anyhow, let me know if you're interested and we can go from there.

Cheers!

MB

Mike Bridger, M.Sc., R.P.Bio.



Caribou Recovery Biologist | Northeast Region

Ministry of Forests, Lands, Natural Resource Operations and Rural Development

Suite 400, 10003-110th Avenue, Fort St. John, BC, V1J 6M7

Ph: 250-787-3294



--

Rob Serrouya

University of Alberta | ABMI | Caribou Monitoring Unit

Resources for Woodland Caribou Science and Recovery:

<https://cmu.abmi.ca/>

Columbia Mountains Caribou Research Project

Box 9158

Revelstoke, British Columbia, V0E 3K0

--

Rob Serrouya

University of Alberta | ABMI | Caribou Monitoring Unit

Resources for Woodland Caribou Science and Recovery:

<https://cmu.abmi.ca/>

Columbia Mountains Caribou Research Project

Box 9158

Revelstoke, British Columbia, V0E 3K0



--

Rob Serrouya

University of Alberta | ABMI | Caribou Monitoring Unit

Resources for Woodland Caribou Science and Recovery:

<https://cmu.abmi.ca/>

Columbia Mountains Caribou Research Project

Box 9158

Revelstoke, British Columbia, V0E 3K0

--

Rob Serrouya

University of Alberta | ABMI | Caribou Monitoring Unit

Resources for Woodland Caribou Science and Recovery:

<https://cmu.abmi.ca/>

Columbia Mountains Caribou Research Project

Box 9158

Revelstoke, British Columbia, V0E 3K0

Page 91 of
92
92-AR-
2019-96488

Page 92 of
92 FNR-
2019-96488